



GB05/545

INVESTOR IN PEOPLE

The Patent Office
 Concept House
 Cardiff Road
 Newport
 South Wales
 NP10 8QQ

REC'D 10 MAR 2005

WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.


Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Dated

20 January 2005

PRIORITY DOCUMENT
 SUBMITTED OR TRANSMITTED IN
 COMPLIANCE WITH
 RULE 17.1(a) OR (b)



Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference 101379-2

2. Patent application number 0413386.4
(The Patent Office will fill in this part) 16 JUN 2004

3. Full name, address and postcode of the or of each applicant (underline all surnames) AstraZeneca AB
SE-151 85 Sodertalje
Sweden

78 2244 8003

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation Sweden

4. Title of the invention
COMPOUNDS

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)
AstraZeneca
Global Intellectual Property
P O Box 272
Mereside, Alderley Park
Macclesfield,
Cheshire SK10 4GR

Patents ADP number (if you know it) 817 970 7001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body.

See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description

100

Claim(s)

5

Abstract

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents
(please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

S. A. Douglas

Date 15/06/2000

12. Name and daytime telephone number of person to contact in the United Kingdom

Shirley Douglas 01625 510057

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

COMPOUNDS

The present invention relates to a group of benzoyl amino heterocyclyl compounds which are useful in the treatment or prevention of a disease or medical condition mediated through glucokinase (GLK or GK), leading to a decreased glucose threshold for insulin secretion. In addition the compounds are predicted to lower blood glucose by increasing hepatic glucose uptake. Such compounds may have utility in the treatment of Type 2 diabetes and obesity. The invention also relates to pharmaceutical compositions comprising said compounds and to methods of treatment of diseases mediated by GLK using said compounds.

In the pancreatic β -cell and liver parenchymal cells the main plasma membrane glucose transporter is GLUT2. Under physiological glucose concentrations the rate at which GLUT2 transports glucose across the membrane is not rate limiting to the overall rate of glucose uptake in these cells. The rate of glucose uptake is limited by the rate of phosphorylation of glucose to glucose-6-phosphate (G-6-P) which is catalysed by glucokinase (GLK) [1]. GLK has a high (6-10mM) K_m for glucose and is not inhibited by physiological concentrations of G-6-P [1]. GLK expression is limited to a few tissues and cell types, most notably pancreatic β -cells and liver cells (hepatocytes) [1]. In these cells GLK activity is rate limiting for glucose utilisation and therefore regulates the extent of glucose induced insulin secretion and hepatic glycogen synthesis. These processes are critical in the maintenance of whole body glucose homeostasis and both are dysfunctional in diabetes [2].

In one sub-type of diabetes, Maturity-Onset Diabetes of the Young Type 2 (MODY-2), the diabetes is caused by GLK loss of function mutations [3, 4]. Hyperglycaemia in MODY-2 patients results from defective glucose utilisation in both the pancreas and liver [5]. Defective glucose utilisation in the pancreas of MODY-2 patients results in a raised threshold for glucose stimulated insulin secretion. Conversely, rare activating mutations of GLK reduce this threshold resulting in familial hyperinsulinism [6, 6a, 7]. In addition to the reduced GLK activity observed in MODY-2 diabetics, hepatic glucokinase activity is also decreased in type 2 diabetics [8]. Importantly, global or liver selective overexpression of GLK prevents or reverses the development of the diabetic phenotype in both dietary and genetic models of the disease [9-12]. Moreover, acute treatment of type 2 diabetics with fructose improves glucose tolerance through stimulation of hepatic glucose utilisation [13]. This effect is believed to be

mediated through a fructose induced increase in cytosolic GLK activity in the hepatocyte by the mechanism described below [13].

Hepatic GLK activity is inhibited through association with GLK regulatory protein (GLKRP). The GLK/GLKRP complex is stabilised by fructose-6-phosphate (F6P) binding to the GLKRP and destabilised by displacement of this sugar phosphate by fructose-1-phosphate (F1P). F1P is generated by fructokinase mediated phosphorylation of dietary fructose. Consequently, GLK/GLKRP complex integrity and hepatic GLK activity is regulated in a nutritionally dependent manner as F6P is dominant in the post-absorptive state whereas F1P predominates in the post-prandial state. In contrast to the hepatocyte, the pancreatic β -cell expresses GLK in the absence of GLKRP. Therefore, β -cell GLK activity is regulated extensively by the availability of its substrate, glucose. Small molecules may activate GLK either directly or through destabilising the GLK/GLKRP complex. The former class of compounds are predicted to stimulate glucose utilisation in both the liver and the pancreas whereas the latter are predicted to act exclusively in the liver. However, compounds with either profile are predicted to be of therapeutic benefit in treating Type 2 diabetes as this disease is characterised by defective glucose utilisation in both tissues.

GLK, GLKRP and the K_{ATP} channel are expressed in neurones of the hypothalamus, a region of the brain that is important in the regulation of energy balance and the control of food intake [14-18]. These neurones have been shown to express orectic and anorectic neuropeptides [15, 19, 20] and have been assumed to be the glucose-sensing neurones within the hypothalamus that are either inhibited or excited by changes in ambient glucose concentrations [17, 19, 21, 22]. The ability of these neurones to sense changes in glucose levels is defective in a variety of genetic and experimentally induced models of obesity [23-28]. Intracerebroventricular (icv) infusion of glucose analogues, that are competitive inhibitors of glucokinase, stimulate food intake in lean rats [29, 30]. In contrast, icv infusion of glucose suppresses feeding [31]. Thus, small molecule activators of GLK may decrease food intake and weight gain through central effects on GLK. Therefore, GLK activators may be of therapeutic use in treating eating disorders, including obesity, in addition to diabetes. The hypothalamic effects will be additive or synergistic to the effects of the same compounds acting in the liver and/or pancreas in normalising glucose homeostasis, for the treatment of Type 2 diabetes. Thus the GLK/GLKRP system can be described as a potential "Diabesity" target (of benefit in both Diabetes and Obesity).


GLK is also expressed in specific entero-endocrine cells where it is believed to control the glucose sensitive secretion of the incretin peptides GIP (glucose-dependent insulintropic polypeptide) and GLP-1 (Glucagon-Like Peptide-1) from gut K-cells and L-cells respectively (32, 33, 34). Therefore, small molecule activators of GLK may have additional beneficial effects on insulin secretion, b-cell function and survival and body weight as a consequence of stimulating GIP and GLP-1 secretion from these entero-endocrine cells.

In WO00/58293 and WO01/44216 (Roche), a series of benzylcarbamoyl compounds are described as glucokinase activators. The mechanism by which such compounds activate GLK is assessed by measuring the direct effect of such compounds in an assay in which GLK activity is linked to NADH production, which in turn is measured optically - see details of the *in vitro* assay described hereinafter. Compounds of the present invention may activate GLK directly or may activate GLK by inhibiting the interaction of GLKRP with GLK.

Further GLK activators have been described in WO03/095438 (substituted phenylacetamides, Roche), WO03/055482 (carboxamide and sulphonamide derivatives, Novo Nordisk), WO2004/002481 (arylcarbonyl derivatives, Novo Nordisk), and in WO03/080585 (amino-substituted benzoylaminoheterocycles, Banyu).

Our International application Number: WO03/000267 describes a group of benzoyl amino pyridyl carboxylic acids which are activators of the enzyme glucokinase (GLK).

Our International application Number: WO03/015774 describes compounds of the Formula (A):




(A)

wherein R^3 is a substituted heterocycle, and wherein the substituents R^1 , R^2 and those on the heterocycle R^3 are selected such that the compounds are overall neutral.

We have surprisingly found a small group of compounds, generally a selected subgroup of those described in WO 03/015774, which have generally superior potency for the GLK enzyme, and more advantageous physical properties, including, for example, higher aqueous solubility, higher permeability, and/or lower plasma protein binding. Consequently, such compounds would be expected to display higher plasma free drug levels and superior *in vivo* efficacy after oral dosing as determined, for example, by activity in Oral Glucose

Tolerance Tests (OGTTs). Therefore this group of compounds would be expected to provide superior oral exposure at a lower dose and thereby be particularly suitable for use in the treatment or prevention of a disease or medical condition mediated through GLK.

Thus, according to the first aspect of the invention there is provided a compound of
5 Formula (I):



wherein:

R^1 is methoxymethyl;

10 R^2 is selected from $-C(O)NR^4R^5$, $-SO_2NR^4R^5$, $-S(O)_pR^4$ and HET-2;

HET-1 is a 5- or 6-membered, C-linked heteroaryl ring containing a nitrogen atom in the 2-position and optionally 1 or 2 further ring heteroatoms independently selected from O, N and S; which ring is optionally substituted on an available carbon atom, or on a ring nitrogen atom provided it is not thereby quaternised, with 1 or 2 substituents independently selected from

15 R^6 ;

HET-2 is a 4-, 5- or 6-membered, C- or N-linked heterocyclyl ring containing 1, 2, 3 or 4 heteroatoms independently selected from O, N and S, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$, and wherein a sulphur atom in the heterocyclic ring may optionally be oxidised to a $S(O)$ or $S(O)_2$ group, which ring is optionally substituted on an available

20 carbon or nitrogen atom by 1 or 2 substituents independently selected from R^7 ;

R^3 is selected from halo, fluoromethyl, difluoromethyl, trifluoromethyl, methyl, methoxy and cyano;

R^4 is selected from hydrogen, (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from HET-2, $-OR^5$, $-SO_2R^5$, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R^7) and $-C(O)NR^5R^5$] and HET-2;

25 R^5 is hydrogen or (1-4C)alkyl;

or R^4 and R^5 together with the nitrogen atom to which they are attached may form a heterocyclyl ring system as defined by HET-3;

R^6 is independently selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkylS(O)p(1-4C)alkyl, amino(1-4C)alkyl, (1-4C)alkylamino(1-4C)alkyl, di(1-4C)alkylamino(1-4C)alkyl and HET-4;

R^7 is selected from $-OR^5$, (1-4C)alkyl, $-C(O)(1-4C)alkyl$, $-C(O)NR^4R^5$, (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl and $-S(O)pR^5$;

HET-3 is an N-linked, 4 to 6 membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 or 2 further heteroatoms (in addition to the linking N atom) independently selected from O, N and S, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$ and wherein a sulphur atom in the ring may optionally be oxidised to a S(O) or S(O)₂

10 group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^8 ; or

HET-3 is an N-linked, 7 membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further heteroatom (in addition to the linking N atom) independently selected from O, S and N, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$

15 group and wherein a sulphur atom in the ring may optionally be oxidised to a S(O) or S(O)₂ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^8 ; or

HET-3 is an 6-10 membered bicyclic saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further nitrogen atom (in addition to the linking N atom) wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$; which ring is optionally substituted on an available carbon or nitrogen atom by 1 substituent selected from R^3 ;

20 R^8 is selected from $-OR^5$, (1-4C)alkyl, $-C(O)(1-4C)alkyl$, $-C(O)NR^4R^5$, (1-4C)alkylamino, di(1-4C)alkylamino, HET-3 (wherein said ring is unsubstituted), (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl and $-S(O)pR^5$;

25 HET-4 is a 5- or 6-membered, C- or N- linked unsubstituted heteroaryl ring containing 1, 2 or 3 ring heteroatoms independently selected from O, N and S;

p is (independently at each occurrence) 0, 1 or 2;

m is 0 or 1;

n is 0, 1 or 2;

30 provided that when m is 0, then n is 1 or 2;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention there is provided a compound of the formula (I), or a salt, pro-drug or solvate thereof, wherein:

HET-3 is an N-linked, 4 to 6 membered, saturated or partially unsaturated heterocyclcyl ring, optionally containing 1 or 2 further heteroatoms (in addition to the linking N atom) independently selected from O, N and S, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$ and wherein a sulphur atom in the ring may optionally be oxidised to a $S(O)$ or $S(O)_2$ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^8 .

It will be understood that when R^4 is $-C(O)NR^5R^5$, each R^5 is independently selected from hydrogen and (1-4C)alkyl, and therefore this definition of R^4 includes (but is not limited to) $-CONH_2$, $-CONHMe$, $-CONMe_2$ and $-CONMeEt$.

10 It will be understood that where a compound of the formula (I) contains more than one HET-2 ring, they may be the same or different.

It will be understood that where a compound of the formula (I) contains more than one group R^4 , they may be the same or different.

15 It will be understood that where a compound of the formula (I) contains more than one group R^5 , they may be the same or different.

It will be understood that where a compound of the formula (I) contains more than one group R^8 , they may be the same or different.

A similar convention applies for all other groups and substituents on a compound of formula (I) as hereinbefore defined.

20 Compounds of Formula (I) may form salts which are within the ambit of the invention. Pharmaceutically acceptable salts are preferred although other salts may be useful in, for example, isolating or purifying compounds.


25 In another aspect, the invention relates to compounds of formula (I) as hereinabove defined or to a pharmaceutically acceptable salt.

In another aspect, the invention relates to compounds of formula (I) as hereinabove defined or to a pro-drug thereof. Suitable examples of pro-drugs of compounds of formula (I) are in-vivo hydrolysable esters of compounds of formula (I). Therefore in another aspect, the invention relates to compounds of formula (I) as hereinabove defined or to an in-vivo hydrolysable ester thereof.

30 In this specification the generic term "alkyl" includes both straight-chain and branched-chain alkyl groups. However references to individual alkyl groups such as "propyl"

are specific for the straight chain version only and references to individual branched-chain alkyl groups such as *t*-butyl are specific for the branched chain version only. For example, "(1-4C)alkyl" includes methyl, ethyl, propyl, isopropyl and *t*-butyl. An analogous convention applies to other generic terms.

- 5 For the avoidance of doubt, reference to the group HET-1 containing a nitrogen in the 2-position, is intended to refer to the 2-position relative to the amide nitrogen atom to which the group is attached. For example, the following structures are encompassed (but not limited to):



10

Suitable examples of HET-1 as a 5- or 6-membered, C-linked heteroaryl ring as hereinbefore defined, include thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl and triazolyl.

- 15 It will be understood that HET-2 can be a saturated, or partially or fully unsaturated ring.

- Suitable examples of HET-2 include azetidiny, furyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, morpholino, morpholinyl, piperidiny, piperazinyl, morpholinyl, 20 thiomorpholinyl, pyrrolyl, pyrrolidinyl, pyrrolidonyl, 2,5-dioxopyrrolidinyl, 1,1-dioxotetrahydrothienyl, 2-oxoimidazolidinyl, 2,4-dioxoimidazolidinyl, 2-oxo-1,3,4-(4-triazolinyl), 2-oxazolidinonyl, 2-oxotetrahydrofuranly, tetrahydrofuranly, tetrahydropyranly, 1,1-dioxothiormorpholino, 1,3-dioxolanyl, 1,2,4-triazolyl, 1,2,3-triazolyl, pyranly, and 4-pyridonyl.


- 25 It will be understood that HET-2 may be linked by any appropriate available C or N atom, therefore for example, for HET-2 as "imidazolyl" includes 1-, 2-, 4- and 5- imidazolyl.

Suitable examples of HET-3 as a 4-6 membered saturated or partially unsaturated heterocyclic ring are morpholino, piperidiny, piperazinyl, pyrrolidinyl and azetidiny.

- Suitable examples of HET-3 as a 7-membered saturated or partially unsaturated 30 heterocyclic ring are homopiperazinyl, homo-morpholino, homo-thiomorpholino (and

versions thereof wherein the sulfur is oxidised to an SO or S(O)₂ group) and homopiperidinyl.

Suitable examples of HET-3 as an 6-10 membered bicyclic heterocyclic ring are bicyclic saturated or partially unsaturated heterocyclyl ring such as those illustrated by the structures shown below (wherein the dotted line indicates the point of attachment to the rest of the molecule):



Suitable examples of HET-4 are furyl, pyrrolyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl and triazolyl.

It will be appreciated that, where definitions of heterocyclyl groups HET-1 to HET-4 encompass heteroaryl rings which may be substituted on nitrogen, such substitution may not result in charged quaternary nitrogen atoms. It will be appreciated that the definitions of HET-1 to HET-4 are not intended to include any O-O, O-S or S-S bonds. It will be appreciated that the definitions of HET-1 to HET-4 are not intended to include unstable structures.

Examples of **(1-4C)alkyl** include methyl, ethyl, propyl, isopropyl, butyl and tert-butyl; examples of **(3-6C)cycloalkyl** include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; examples of **halo** include fluoro, chloro, bromo and iodo; examples of **hydroxy(1-4C)alkyl** include hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 5 1-hydroxyisopropyl and 4-hydroxybutyl; examples of **(1-4C)alkoxy(1-4C)alkyl** include methoxymethyl, ethoxymethyl, tert-butoxymethyl, 2-methoxyethyl, 2-ethoxyethyl, methoxypropyl, 2-methoxypropyl and methoxybutyl; examples of **(1-4C)alkylS(O)p(1-4C)alkyl** include methylsulfinylmethyl, ethylsulfinylmethyl, ethylsulfinylethyl, methylsulfinylpropyl, methylsulfinylbutyl, methylsulfonylmethyl, ethylsulfonylmethyl, 10 ethylsulfonylethyl, methylsulfonylpropyl, methylsulfonylbutyl, methylthiomethyl, ethylthiomethyl, ethylthioethyl, methylthiopropyl, and methylthiobutyl; examples of **amino(1-4C)alkyl** include aminomethyl, aminoethyl, 2-aminopropyl, 3-aminopropyl, 1-aminoisopropyl and 4-aminobutyl; examples of **(1-4C)alkylamino(1-4C)alkyl** include (N-methyl)aminomethyl, (N-ethyl)aminomethyl, 1-((N-methyl)amino)ethyl, 2-((N-methyl)amino)ethyl, (N-ethyl)aminoethyl, (N-methyl)aminopropyl, and 4-((N-methyl)amino)butyl; examples of **di(1-4C)alkylamino(1-4C)alkyl** include dimethylaminomethyl, methyl(ethyl)aminomethyl, methyl(ethyl)aminoethyl, (N,N-diethyl)aminoethyl, (N,N-dimethyl)aminopropyl and (N,N-dimethyl)aminobutyl; examples of **(1-4C)alkylamino** include methylamino, ethylamino, propylamino, isopropylamino, 20 butylamino and tert-butylamino; examples of **di(1-4C)alkylamino** include dimethylamino, methyl(ethyl)amino, diethylamino, dipropylamino, di-isopropylamino and dibutylamino; examples of **-C(O)(1-4C)alkyl** include methylcarbonyl, ethylcarbonyl, propylcarbonyl and tert-butyl carbonyl.


25 It is to be understood that, insofar as certain of the compounds of Formula (I) defined above may exist in optically active or racemic forms by virtue of one or more asymmetric carbon atoms, the invention includes in its definition any such optically active or racemic form which possesses the property of stimulating GLK directly or inhibiting the GLK/GLKRP interaction. The synthesis of optically active forms may be carried out by standard techniques 30 of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form. It is also to be understood that certain

compounds may exist in tautomeric forms and that the invention also relates to any and all tautomeric forms of the compounds of the invention which activate GLK.

In one embodiment of the invention are provided compounds of formula (I), in an alternative embodiment are provided pharmaceutically-acceptable salts of compounds of formula (I), in a further alternative embodiment are provided in-vivo hydrolysable esters of compounds of formula (I), and in a further alternative embodiment are provided pharmaceutically-acceptable salts of in-vivo hydrolysable esters of compounds of formula (I).

Preferred values of each variable group are as follows. Such values may be used where appropriate with any of the values, definitions, claims, aspects or embodiments defined hereinbefore or hereinafter.

(1) R^1 is methoxymethyl and the configuration is preferably (S), that is:



(2) R^2 is $-C(O)NR^4R^5$

(3) R^2 is $-SO_2NR^4R^5$

15 (4) R^2 is $-S(O)_pR^4$

(5) R^2 is HET-2

(6) m is 1 and R^2 is in the para position relative to the ether linkage

(7) m is 1 and n is 0 or 1

(8) m is 1 and n is 0

20 (9) m is 1, n is 0 and R^2 is in the para position relative to the ether linkage

(10) n is 0

(11) n is 1

(12) n is 2

(13) n is 2 and both R^3 are halo

25 (14) R^3 is halo or trifluoromethyl

(15) R^3 is halo

(16) R^3 is chloro or fluoro

(17) R^3 is fluoro

(18) n is 2 and both R^3 are fluoro,

30 (19) n is 2, both R^3 are fluoro and are in the 3- and 5-positions relative to the ether linkage

- (20) p is 0
- (21) p is 1
- (22) p is 2
- (23) HET-1 is a 5-membered heteroaryl ring
- 5 (24) HET-1 is a 6-membered heteroaryl ring
- (25) HET-1 is substituted with 1 or 2 substituents independently selected from R⁶
- (26) HET-1 is substituted with 1 substituent selected from R⁶
- (27) HET-1 is unsubstituted
- (28) HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl,
- 10 pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, and triazolyl
- (29) HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl and oxadiazolyl
- (30) HET-1 is selected from pyridyl, pyrazinyl, pyridazinyl and pyrimidinyl
- 15 (31) HET-1 is selected from thiazolyl, pyrazolyl and oxazolyl
- (32) HET-1 is selected from thiadiazolyl and oxadiazolyl
- (33) HET-1 is selected from 1,3,4-thiadiazolyl and 1,3,4-oxadiazolyl
- (34) HET-1 is selected from 1,2,4-oxadiazolyl and 1,2,4-oxadiazolyl
- (35) HET-1 is pyridyl
- 20 (36) HET-1 is pyrazinyl
- (37) R⁶ is selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, di(1-4C)alkylamino(1-4C)alkyl and HET-4
- (38) R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, hydroxymethyl, methoxymethyl, aminomethyl, N-methylaminomethyl, dimethylaminomethyl
- 25 (39) R⁶ is selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkylS(O)p(1-4C)alkyl, amino(1-4C)alkyl, (1-4C)alkylamino(1-4C)alkyl, and di(1-4C)alkylamino(1-4C)alkyl
- (40) R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl
- 30 (41) R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, hydroxymethyl and methoxymethyl
- (42) R⁶ is selected from methyl, ethyl, bromo, chloro and fluoro
- (43) R⁶ is methyl

- (44) R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, dimethylaminomethyl, hydroxymethyl and methoxymethyl
- (45) when 2 substituents R⁶ are present, both are selected from methyl, ethyl, bromo, chloro and fluoro, preferably both are methyl
- 5 (46) HET-4 is selected from furyl, pyrrolyl and thienyl
- (47) HET-4 is furyl
- (48) R⁴ is hydrogen
- (49) R⁴ is (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from HET-2, -OR⁵, -SO₂R⁵, (3-6C)cycloalkyl (optionally substituted with 1 group selected
- 10 from R⁷) and -C(O)NR⁵R⁵]
- (50) R⁴ is (1-4C)alkyl [optionally substituted by 1 substituent selected from HET-2, -OR⁵, -SO₂R⁵, (3-6C)cycloalkyl and -C(O)NR⁵R⁵]
- (51) R⁴ is HET-2
- (52) R⁴ is selected from hydrogen, (1-4C)alkyl, and (1-4C)alkyl substituted with -OR⁵
- 15 (53) HET-2 is unsubstituted
- (54) HET-2 is substituted with 1 or 2 substituents independently selected from (1-4C)alkyl, hydroxy and (1-4C)alkoxy
- (55) HET-2 is a fully saturated ring system
- (56) HET-2 is a fully unsaturated ring system
- 20 (57) HET-2 is selected from azetidiny, morpholino, morpholinyl, piperidiny, piperazinyl, 3-oxopiperazinyl, thiomorpholinyl, pyrrolidinyl, pyrrolidonyl, 2,5-dioxopyrrolidinyl, 1,1-dioxotetrahydrothienyl, 2-oxazolidinonyl, 2-oxotetrahydrofuranyl, tetrahydrofuranyl, tetrahydropyranyl, 1,1-dioxothiomorpholino, 1,3-dioxolanyl, 2-oxoimidazolidinyl, 2,4-dioxoimidazolidinyl, pyranly and 4-pyridonyl
- 25 (58) HET-2 is selected from azetidiny, morpholino, morpholinyl, piperidiny, piperazinyl, pyrrolidinyl, thiomorpholinyl, tetrahydrofuranyl, and tetrahydropyranyl
- (59) HET-2 is selected from furyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, pyrrolyl, 1,2,4-triazolyl and 1,2,3-triazolyl
- 30 (60) HET-2 is selected from furyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, piperidiny, piperazinyl, 3-oxopiperazinyl, pyrrolidinyl, pyrrolidonyl, 2-oxazolidinonyl, tetrahydrofuranyl, tetrahydropyranyl, 1,1-dioxotetrahydrothienyl, and 2-oxoimidazolidinyl

(61) HET-2 is selected from morpholino, furyl, imidazolyl, oxazolyl, isoxazolyl, oxadiazolyl, piperidinyl, piperazinyl, 3-oxopiperazinyl, pyrrolidinyl, 2-pyrrolidonyl, 2-oxazolidinonyl, tetrahydrofuranyl, tetrahydropyranyl, 1,1-dioxotetrahydrothienyl, and 2-oxoimidazolidinyl

(62) HET-2 is selected from morpholino, furyl, imidazolyl, isoxazolyl, oxadiazolyl,

5 piperidinyl, piperazinyl, 3-oxopiperazinyl, pyrrolidinyl, 2-pyrrolidonyl, tetrahydropyranyl, 1,1-dioxotetrahydrothienyl, and 2-oxoimidazolidinyl

(63) R^5 is hydrogen

(64) R^5 is (1-4)alkyl

(65) R^5 is hydrogen or methyl

10 (66) R^7 is selected from $-OR^5$, (1-4C)alkyl, $-C(O)(1-4C)alkyl$, $-C(O)NR^4R^5$, (1-4C)alkoxy(1-4C)alkyl, and hydroxy(1-4C)alkyl

(67) R^7 is selected from $-OR^5$, (1-4C)alkyl, $-C(O)(1-4C)alkyl$, $-C(O)NR^4R^5$, and hydroxy(1-4C)alkyl

(68) R^7 is selected from hydroxy, methoxy, $-COMe$, $-CONH_2$, $-CONHMe$, $-CONMe_2$, and

15 hydroxymethyl

(69) R^7 is selected from (1-4C)alkyl, hydroxy and (1-4C)alkoxy

(70) R^7 is selected from methyl, ethyl, methoxy and hydroxy

(71) R^8 is selected from methyl, hydroxy, methoxy, $-COMe$, $-CONH_2$, $-CONHMe$, $-CONMe_2$,

hydroxymethyl, hydroxyethyl, $-NHMe$ and $-NMe_2$ (72) R^8 is selected from morpholino,

20 piperidinyl, piperazinyl, pyrrolidinyl and azetidiny

(73) R^8 is selected from methyl, $-COMe$, $-CONH_2$, hydroxyethyl and hydroxy

(74) HET-3 is a fully saturated ring

(75) HET-3 is selected from morpholino, piperidinyl, piperazinyl, pyrrolidinyl and azetidiny

(76) R^4 and R^5 together with the nitrogen to which they are attached form a ring as defined by

25 HET-3

(77) HET-3 is a 4 to 6-membered saturated or partially unsaturated heterocyclic ring

(78) HET-3 is a 7-membered saturated or partially unsaturated heterocyclic ring

(79) HET-3 is an 6 to 10-membered bicyclic saturated or partially unsaturated heterocyclic ring

30

According to a further feature of the invention there is provided the following preferred groups of compounds of the invention:

In a further aspect of the invention there is provided a compound of Formula (I)

wherein:

R¹ is methoxymethyl;

R² is selected from -C(O)NR⁴R⁵, -SO₂NR⁴R⁵, -S(O)_pR⁴ and HET-2;

HET-1 is a 5- or 6-membered, C-linked heteroaryl ring containing a nitrogen atom in the 2-
5 position and optionally 1, 2 or 3 further ring heteroatoms independently selected from O, N
and S; which ring is optionally substituted on an available carbon atom, or on a ring nitrogen
atom provided it is not thereby quaternised, with 1 or 2 substituents independently selected
from R⁶;

HET-2 is a 5- or 6-membered, C- or N-linked heterocyclyl ring containing 1, 2, 3 or 4
10 heteroatoms independently selected from O, N and S, wherein a -CH₂- group can optionally
be replaced by a -C(O)-, and wherein a sulphur atom in the heterocyclic ring may optionally
be oxidised to an S(O) or S(O)₂ group, which ring is optionally substituted on an available
carbon or nitrogen atom by 1 or 2 substituents independently selected from R⁷;

R³ is selected from halo, fluoromethyl, difluoromethyl, trifluoromethyl, methyl, methoxy and
15 cyano;

R⁴ is selected from hydrogen, (1-4C)alkyl, [optionally substituted by -OR⁵] and HET-2;

R⁵ is hydrogen or (1-4C)alkyl;

or R⁴ and R⁵ together with the nitrogen atom to which they are attached may form a 4-6
membered heterocyclyl ring system as defined by HET-3;

20 R⁶ is independently selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, (1-4C)alkoxy(1-
4C)alkyl, (1-4C)alkylS(O)_p(1-4C)alkyl, amino(1-4C)alkyl, (1-4C)alkylamino(1-4C)alkyl,
di(1-4C)alkylamino(1-4C)alkyl and HET-4;

R⁷ is selected from -OR⁵ and (1-4C)alkyl;

HET-3 is an N-linked, 4 to 6 membered, saturated or partially unsaturated heterocyclyl ring,
25 optionally containing 1 or 2 further heteroatoms (in addition to the linking N atom)
independently selected from O, N and S, wherein a -CH₂- group can optionally be replaced by
a -C(O)- and wherein a sulphur atom in the ring may optionally be oxidised to an S(O) or
S(O)₂ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1
or 2 substituents independently selected from R⁸;

30 R⁸ is selected from -OR⁵ and (1-4C)alkyl;

HET-4 is a 5- or 6-membered, C- or N- linked unsubstituted heteroaryl ring containing 1, 2 or
3 ring heteroatoms independently selected from O, N and S;

p is (independently at each occurrence) 0, 1 or 2;

m is 0 or 1;

n is 0, 1 or 2;

provided that when m is 0, then n is 1 or 2;

5 or a salt, pro-drug or solvate thereof.

In a further aspect of the invention there is provided a compound of Formula (I)

wherein:

R¹ is methoxymethyl;

10 R² is selected from -C(O)NR⁴R⁵, -SO₂NR⁴R⁵, -S(O)_pR⁴ and HET-2;

HET-1 is a 5- or 6-membered, C-linked heteroaryl ring containing a nitrogen atom in the 2-position and optionally 1, 2 or 3 further ring heteroatoms independently selected from O, N and S; which ring is optionally substituted on an available carbon atom, or on a ring nitrogen atom provided it is not thereby quaternised, with 1 or 2 substituents independently selected

15 from R⁶;

HET-2 is a 5- or 6-membered, C- or N-linked heterocyclyl ring containing 1, 2, 3 or 4 heteroatoms independently selected from O, N and S, wherein a -CH₂- group can optionally be replaced by a -C(O)-, and wherein a sulphur atom in the heterocyclic ring may optionally be oxidised to an S(O) or S(O)₂ group, which ring is optionally substituted on an available

20 carbon or nitrogen atom by 1 or 2 substituents independently selected from R⁷;

R³ is selected from halo, fluoromethyl, difluoromethyl, trifluoromethyl, methyl, methoxy and cyano;

R⁴ is selected from hydrogen, (1-4C)alkyl, [optionally substituted by -OR⁵] and HET-2;

R⁵ is hydrogen or (1-4C)alkyl;

25 or R⁴ and R⁵ together with the nitrogen atom to which they are attached may form a heterocyclyl ring system as defined by HET-3;

R⁶ is independently selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkylS(O)_p(1-4C)alkyl, amino(1-4C)alkyl, (1-4C)alkylamino(1-4C)alkyl, di(1-4C)alkylamino(1-4C)alkyl and HET-4;

30 R⁷ is selected from -OR⁵ and (1-4C)alkyl;

HET-3 is an N-linked, 4 to 6 membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 or 2 further heteroatoms (in addition to the linking N atom)

independently selected from O, N and S, wherein a $-\text{CH}_2-$ group can optionally be replaced by a $-\text{C}(\text{O})-$ and wherein a sulphur atom in the ring may optionally be oxidised to an $\text{S}(\text{O})$ or $\text{S}(\text{O})_2$ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^8 ; or

- 5 HET-3 is an N-linked, 7 membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further heteroatom (in addition to the linking N atom) independently selected from O, S and N, wherein a $-\text{CH}_2-$ group can optionally be replaced by a $-\text{C}(\text{O})-$ group and wherein a sulphur atom in the ring may optionally be oxidised to an $\text{S}(\text{O})$ or $\text{S}(\text{O})_2$ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2
- 10 substituents independently selected from R^8 ; or
- HET-3 is an 6-10 membered bicyclic saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further nitrogen atom (in addition to the linking N atom) wherein a $-\text{CH}_2-$ group can optionally be replaced by a $-\text{C}(\text{O})-$; which ring is optionally substituted on an available carbon or nitrogen atom by 1 substituent selected from R^3 ;
- 15 R^8 is selected from $-\text{OR}^5$ and (1-4C)alkyl;
- HET-4 is a 5- or 6-membered, C- or N- linked unsubstituted heteroaryl ring containing 1, 2 or 3 ring heteroatoms independently selected from O, N and S;
- p is (independently at each occurrence) 0, 1 or 2;
- m is 0 or 1;
- 20 n is 0, 1 or 2;
- provided that when m is 0, then n is 1 or 2;
- or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as

- 25 hereinbefore defined wherein:

R^1 is methoxymethyl;

R^2 is selected from $-\text{C}(\text{O})\text{NR}^4\text{R}^5$, $-\text{SO}_2\text{NR}^4\text{R}^5$, $-\text{S}(\text{O})_p\text{R}^4$ and HET-2;

HET-1 is a 5- or 6-membered, C-linked heteroaryl ring containing a nitrogen atom in the 2-position and optionally 1 or 2 further ring heteroatoms independently selected from O, N and

- 30 S; which ring is optionally substituted on an available carbon atom, or on a ring nitrogen atom provided it is not thereby quaternised, with 1 or 2 substituents independently selected from R^6 ;

HET-2 is a 4-, 5- or 6-membered, C- or N-linked heterocyclyl ring containing 1, 2, 3 or 4 heteroatoms independently selected from O, N and S, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$, and wherein a sulphur atom in the heterocyclic ring may optionally be oxidised to an $S(O)$ or $S(O)_2$ group, which ring is optionally substituted on an available

5 carbon or nitrogen atom by 1 or 2 substituents independently selected from R^7 ;

R^3 is selected from halo, fluoromethyl, difluoromethyl, trifluoromethyl, methyl, methoxy and cyano;

R^4 is selected from (1-4C)alkyl [substituted by 1 or 2 substituents independently selected from HET-2, $-SO_2R^5$, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R^7) and

10 $-C(O)NR^5R^5$];

R^5 is hydrogen or (1-4C)alkyl;

or R^4 and R^5 together with the nitrogen atom to which they are attached may form a 4-6 membered heterocyclyl ring system as defined by HET-3;

R^6 is independently selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, (1-4C)alkoxy(1-

15 4C)alkyl, (1-4C)alkyl $S(O)p$ (1-4C)alkyl, amino(1-4C)alkyl, (1-4C)alkylamino(1-4C)alkyl, di(1-4C)alkylamino(1-4C)alkyl and HET-4;

R^7 is selected from $-C(O)(1-4C)alkyl$, $-C(O)NR^4R^5$, (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl and $-S(O)pR^5$;

HET-3 is an N-linked, 4 to 6 membered, saturated or partially unsaturated heterocyclyl ring,

20 optionally containing 1 or 2 further heteroatoms (in addition to the linking N atom)

independently selected from O, N and S, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$ and wherein a sulphur atom in the ring may optionally be oxidised to an $S(O)$ or $S(O)_2$ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^8 ;

25 R^8 is selected from $-C(O)(1-4C)alkyl$, $-C(O)NR^4R^5$, (1-4C)alkylamino, di(1-4C)alkylamino, HET-3 (wherein said ring is unsubstituted), (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl and $-S(O)pR^5$;

HET-4 is a 5- or 6-membered, C- or N- linked unsubstituted heteroaryl ring containing 1, 2 or 3 ring heteroatoms independently selected from O, N and S;

30 p is (independently at each occurrence) 0, 1 or 2;

m is 0 or 1;

n is 0, 1 or 2;

provided that when m is 0, then n is 1 or 2;
or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as

5 hereinbefore defined wherein:

R^1 is methoxymethyl;

R^2 is selected from $-C(O)NR^4R^5$, $-SO_2NR^4R^5$, $-S(O)_pR^4$ and HET-2;

HET-1 is a 5- or 6-membered, C-linked heteroaryl ring containing a nitrogen atom in the 2-position and optionally 1 or 2 further ring heteroatoms independently selected from O, N and
10 S; which ring is optionally substituted on an available carbon atom, or on a ring nitrogen atom provided it is not thereby quaternised, with 1 or 2 substituents independently selected from R^6 ;

HET-2 is a 4-, 5- or 6-membered, C- or N-linked heterocyclyl ring containing 1, 2, 3 or 4 heteroatoms independently selected from O, N and S, wherein a $-CH_2-$ group can optionally
15 be replaced by a $-C(O)-$, and wherein a sulphur atom in the heterocyclic ring may optionally be oxidised to an S(O) or S(O)₂ group, which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^7 ;

R^3 is selected from halo, fluoromethyl, difluoromethyl, trifluoromethyl, methyl, methoxy and cyano;

20 R^4 is selected from (1-4C)alkyl [substituted by 1 or 2 substituents independently selected from HET-2, $-SO_2R^5$, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R^7) and $-C(O)NR^5R^5$];

R^5 is hydrogen or (1-4C)alkyl;

or R^4 and R^5 together with the nitrogen atom to which they are attached may form a

25 heterocyclyl ring system as defined by HET-3;

R^6 is independently selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkylS(O)_p(1-4C)alkyl, amino(1-4C)alkyl, (1-4C)alkylamino(1-4C)alkyl, di(1-4C)alkylamino(1-4C)alkyl and HET-4;

R^7 is selected from $-C(O)(1-4C)alkyl$, $-C(O)NR^4R^5$, (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-
30 4C)alkyl and $-S(O)_pR^5$;

HET-3 is an N-linked, 4 to 6 membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 or 2 further heteroatoms (in addition to the linking N atom)

independently selected from O, N and S, wherein a $-\text{CH}_2-$ group can optionally be replaced by a $-\text{C}(\text{O})-$ and wherein a sulphur atom in the ring may optionally be oxidised to an $\text{S}(\text{O})$ or $\text{S}(\text{O})_2$ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^8 ; or

5 HET-3 is an N-linked, 7 membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further heteroatom (in addition to the linking N atom) independently selected from O, S and N, wherein a $-\text{CH}_2-$ group can optionally be replaced by a $-\text{C}(\text{O})-$ group and wherein a sulphur atom in the ring may optionally be oxidised to an $\text{S}(\text{O})$ or $\text{S}(\text{O})_2$ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2

10 substituents independently selected from R^8 ; or

HET-3 is an 6-10 membered bicyclic saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further nitrogen atom (in addition to the linking N atom), wherein a $-\text{CH}_2-$ group can optionally be replaced by a $-\text{C}(\text{O})-$; which ring is optionally substituted on an available carbon or nitrogen atom by 1 substituent selected from R^3 ;

15 R^8 is selected from $-\text{C}(\text{O})(1-4\text{C})\text{alkyl}$, $-\text{C}(\text{O})\text{NR}^4\text{R}^5$, $(1-4\text{C})\text{alkylamino}$, $\text{di}(1-4\text{C})\text{alkylamino}$, HET-3 (wherein said ring is unsubstituted), $(1-4\text{C})\text{alkoxy}(1-4\text{C})\text{alkyl}$, $\text{hydroxy}(1-4\text{C})\text{alkyl}$ and $-\text{S}(\text{O})\text{pR}^5$;

HET-4 is a 5- or 6-membered, C- or N- linked unsubstituted heteroaryl ring containing 1, 2 or 3 ring heteroatoms independently selected from O, N and S;

20 p is (independently at each occurrence) 0, 1 or 2;

m is 0 or 1;

n is 0, 1 or 2;

provided that when m is 0, then n is 1 or 2;

or a salt, pro-drug or solvate thereof.

25

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0 or 1;

30 HET-1 is a 5- or 6-membered heteroaryl ring;

R^2 is $-\text{CONR}^4\text{R}^5$ or $-\text{SO}_2\text{NR}^4\text{R}^5$;

R^3 is halo or trifluoromethyl;

R^4 is (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from HET-2, $-OR^5$, $-SO_2R^5$, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R^7) and $-C(O)NR^5R^5$];

R^5 is hydrogen or methyl;

5 HET-2 is a 5- or 6- membered heterocyclyl ring as hereinbefore defined, containing 1 or 2 heteroatoms independently selected from O, N and S; and

R^7 is selected from $-OR^5$ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

10 In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is a 5- or 6-membered heteroaryl ring;

15 R^2 is $-CONR^4R^5$ or $-SO_2NR^4R^5$;

R^3 is halo or trifluoromethyl;

R^4 is (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from HET-2, $-OR^5$, $-SO_2R^5$, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R^7) and $-C(O)NR^5R^5$];

20 R^5 is hydrogen or methyl;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is a 5- or 6- membered heterocyclyl ring as hereinbefore defined, containing 1 or 2 heteroatoms independently selected from O, N and S; and

25 R^7 is selected from $-OR^5$ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

30 R^1 is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

R^2 is $-\text{CONR}^4\text{R}^5$ or $-\text{SO}_2\text{NR}^4\text{R}^5$;

R^3 is halo or trifluoromethyl;

R^4 is (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from HET-2, $-\text{OR}^5$, $-\text{SO}_2\text{R}^5$, (3-6C)cycloalkyl and $-\text{C}(\text{O})\text{NR}^5\text{R}^5$];

5 R^5 is hydrogen or methyl;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is selected from azetidiny, morpholino, morpholinyl, piperidiny, piperaziny, 3-oxopiperaziny, thiomorpholinyl, pyrrolidiny, pyrrolidonyl, 2,5-dioxopyrrolidiny, 1,1-

10 dioxotetrahydrothienyl, 2-oxazolidinonyl, 2-oxotetrahydrofuranyl, tetrahydrofuranyl, tetrahydropyranyl, 1,1-dioxothiomorpholino, 1,3-dioxolanyl, 2-oxoimidazolidiny, 2,4-dioxoimidazolidiny, pyranly and 4-pyridonyl; and

R^7 is selected from $-\text{OR}^5$ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

15

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0 or 1;

20 HET-1 is selected from pyridyl, pyrazinyl, pyridaziny and pyrimidinyl;

R^2 is $-\text{CONR}^4\text{R}^5$ or $-\text{SO}_2\text{NR}^4\text{R}^5$;

R^3 is halo or trifluoromethyl;

R^4 is (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from HET-2, $-\text{OR}^5$, $-\text{SO}_2\text{R}^5$, (3-6C)cycloalkyl and $-\text{C}(\text{O})\text{NR}^5\text{R}^5$];

25 R^5 is hydrogen or methyl;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is selected from azetidiny, morpholino, morpholinyl, piperidiny, piperaziny, 3-oxopiperaziny, thiomorpholinyl, pyrrolidiny, pyrrolidonyl, 2,5-dioxopyrrolidiny, 1,1-

30 dioxotetrahydrothienyl, 2-oxazolidinonyl, 2-oxotetrahydrofuranyl, tetrahydrofuranyl, tetrahydropyranyl, 1,1-dioxothiomorpholino, 1,3-dioxolanyl, 2-oxoimidazolidiny, 2,4-dioxoimidazolidiny, pyranly and 4-pyridonyl; and

R^7 is selected from $-\text{OR}^5$ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

5 R^1 is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

R^2 is $-\text{CONR}^4\text{R}^5$ or $-\text{SO}_2\text{NR}^4\text{R}^5$;

10 R^3 is halo or trifluoromethyl;

R^4 is (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from HET-2, $-\text{OR}^5$, $-\text{SO}_2\text{R}^5$, (3-6C)cycloalkyl and $-\text{C}(\text{O})\text{NR}^5\text{R}^5$];

R^5 is hydrogen or methyl;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl,

15 and dimethylaminomethyl;

HET-2 is selected from furyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, pyrrolyl, 1,2,4-triazolyl and 1,2,3-triazolyl; and

R^7 is selected from $-\text{OR}^5$ and (1-4C)alkyl;

20 or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

25 m is 1 and n is 0 or 1;

HET-1 is selected from pyridyl, pyrazinyl, pyridazinyl and pyrimidinyl;

R^2 is $-\text{CONR}^4\text{R}^5$ or $-\text{SO}_2\text{NR}^4\text{R}^5$;

R^3 is halo or trifluoromethyl;

R^4 is (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from

30 HET-2, $-\text{OR}^5$, $-\text{SO}_2\text{R}^5$, (3-6C)cycloalkyl and $-\text{C}(\text{O})\text{NR}^5\text{R}^5$];

R^5 is hydrogen or methyl;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is selected from furyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, pyrrolyl, 1,2,4-triazolyl and 1,2,3-triazolyl; and

R⁷ is selected from -OR⁵ and (1-4C)alkyl;

5 or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

10 m is 1 and n is 0 or 1;

HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

R² is -CONR⁴R⁵ or -SO₂NR⁴R⁵;

R³ is halo or trifluoromethyl;

15 R⁴ is selected from hydrogen, (1-4C)alkyl, [optionally substituted by -OR⁵] and HET-2;

R⁵ is hydrogen or methyl;

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is selected from morpholino, furyl, imidazolyl, isoxazolyl, oxadiazolyl, piperidinyl,

20 piperazinyl, 3-oxopiperazinyl, pyrrolidinyl, 2-pyrrolidonyl, tetrahydropyranyl, 1,1-dioxotetrahydrothienyl, and 2-oxoimidazolidinyl; and

R⁷ is selected from -OR⁵ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

25 In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from pyridyl and pyridazinyl;

30 R² is -CONR⁴R⁵ or -SO₂NR⁴R⁵;

R³ is halo or trifluoromethyl;

R⁴ is selected from hydrogen, (1-4C)alkyl, [optionally substituted by -OR⁵] and HET-2;

R⁵ is hydrogen or methyl;

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is selected from morpholino, furyl, imidazolyl, isoxazolyl, oxadiazolyl, piperidinyl, piperazinyl, 3-oxopiperazinyl, pyrrolidinyl, 2-pyrrolidonyl, tetrahydropyranyl, 1,1-

5 dioxotetrahydrothienyl, and 2-oxoimidazolidinyl; and

R⁷ is selected from -OR⁵ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as

10 hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

15 R² is -CONR⁴R⁵ or -SO₂NR⁴R⁵;

R³ is halo or trifluoromethyl;

R⁴ is selected from (1-4C)alkyl, [optionally substituted by -OR⁵] and HET-2;

R⁵ is hydrogen or methyl;

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl,

20 and dimethylaminomethyl;

HET-2 is selected from piperidinyl, piperazinyl, 3-oxopiperazinyl, 2-pyrrolidonyl, 2,5-dioxopyrrolidinyl, 2-oxotetrahydrofuranlyl, tetrahydrofuranlyl, tetrahydropyranyl, 2-oxoimidazolidinyl, and 2,4-dioxoimidazolidinyl; and

R⁷ is (1-4C)alkyl;

25 or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

30 m is 1 and n is 0 or 1;

HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

R² is -CONR⁴R⁵ or -SO₂NR⁴R⁵;

R^3 is halo or trifluoromethyl;

R^4 is selected from (1-4C)alkyl, [optionally substituted by $-OR^5$] and HET-2;

R^5 is hydrogen or methyl;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl,

5 and dimethylaminomethyl;

HET-2 is piperidiny1 or piperazinyl; and

R^7 is (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

10 In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0 ;

HET-1 is selected from thiazolyl, thiadiazolyl and pyrazolyl;

15 R^2 is $-\text{CONR}^4\text{R}^5$;

R^4 is piperidiny1, optionally substituted with methyl;

R^5 is hydrogen or methyl;

R^6 is methyl;

or a salt, pro-drug or solvate thereof.

20

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0 or 1;

25 HET-1 is selected from pyridyl and pyridazinyl;

R^2 is $-\text{CONR}^4\text{R}^5$ or $-\text{SO}_2\text{NR}^4\text{R}^5$;

R^3 is halo or trifluoromethyl;

R^4 is selected from (1-4C)alkyl, [optionally substituted by $-OR^5$] and HET-2;

R^5 is hydrogen or methyl;

30 R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is selected from piperidinyl, piperazinyl, 3-oxopiperazinyl, 2-pyrrolidinonyl, 2,5-dioxopyrrolidinyl, 2-oxazolidinonyl, 2-oxotetrahydrofuranlyl, tetrahydrofuranlyl, tetrahydropyranlyl, 2-oxoimidazolidinyl, and 2,4-dioxoimidazolidinyl; and

R⁷ is (1-4C)alkyl;

5 or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

10 m is 1 and n is 0 or 1;

HET-1 is selected from pyridyl and pyridazinyl;

R² is -CONR⁴R⁵ or -SO₂NR⁴R⁵;

R³ is halo or trifluoromethyl;

R⁴ is selected from (1-4C)alkyl, [optionally substituted by -OR⁵] and HET-2;

15 R⁵ is hydrogen or methyl;

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is piperidinyl or piperazinyl; and

R⁷ is (1-4C)alkyl;

20 or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

25 HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

R² is -CONR⁴R⁵ or -SO₂NR⁴R⁵;

R³ is halo or trifluoromethyl;

R⁴ and R⁵ together with the nitrogen to which they are attached form a morpholino,

30 piperidinyl, piperazinyl, pyrrolidinyl or azetidiny ring, which ring is optionally substituted on a carbon or nitrogen atom by R⁸;

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl; and

R⁸ is selected from hydroxy, (1-4C)alkoxy and (1-4C)alkyl;
or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as

5 hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, oxazolyl, isoxazolyl
and oxadiazolyl;

10 R² is -CONR⁴R⁵ or -SO₂NR⁴R⁵;

R³ is halo or trifluoromethyl;

R⁴ and R⁵ together with the nitrogen to which they are attached form a morpholino,
piperidinyl, piperazinyl, pyrrolidinyl or azetidiny ring, which ring is optionally substituted on
a carbon or nitrogen atom by R⁸;

15 R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl,
and dimethylaminomethyl; and

R⁸ is pyrrolidine or piperidine;

or a salt, pro-drug or solvate thereof.

20 In a further aspect of the invention is provided a compound of the formula (I) as
hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, oxazolyl, isoxazolyl

25 and oxadiazolyl;

R² is -CONR⁴R⁵ or -SO₂NR⁴R⁵;

R³ is halo or trifluoromethyl;

R⁴ and R⁵ together with the nitrogen to which they are attached form a morpholino,
piperidinyl, piperazinyl, pyrrolidinyl or azetidiny ring, which ring is optionally substituted on

30 a carbon or nitrogen atom by (1-4C)alkyl; and

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl,
and dimethylaminomethyl;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

5 m is 1 and n is 0 or 1;

HET-1 is selected from pyridyl and pyridazinyl;

R^2 is $-\text{CONR}^4\text{R}^5$ or $-\text{SO}_2\text{NR}^4\text{R}^5$;

R^3 is halo or trifluoromethyl;

R^4 and R^5 together with the nitrogen to which they are attached form a morpholino,

10 piperidiny, piperazinyl, pyrrolidiny or azetidiny ring, which ring is optionally substituted on a carbon or nitrogen atom by (1-4C)alkyl; and

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

or a salt, pro-drug or solvate thereof.

15

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0;

20 HET-1 is selected from thiazolyl, thiadiazolyl and pyrazolyl;

R^2 is $-\text{CONR}^4\text{R}^5$;

R^4 and R^5 together with the nitrogen to which they are attached form a piperidiny, or piperazinyl ring, which ring is optionally substituted on a carbon or nitrogen atom by (1-4C)alkyl or by a pyrrolidiny ring;

25 R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as
30 hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0;

HET-1 is selected from thiazolyl, thiadiazolyl and pyrazolyl;

R^2 is $-\text{CONR}^4\text{R}^5$;

R^4 and R^5 together with the nitrogen to which they are attached form an azetidiny ring which ring is optionally substituted on a carbon atom by hydroxy;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, hydroxymethyl, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;
5 or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

10 R^1 is methoxymethyl;

m is 1 and n is 0;

HET-1 is selected from thiazolyl, thiadiazolyl and pyrazolyl;

R^2 is $-\text{CONR}^4\text{R}^5$;

R^4 and R^5 together with the nitrogen to which they are attached form a 7-membered ring HET-

15 3 which ring is optionally substituted on a carbon or nitrogen atom by methyl;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, hydroxymethyl, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;
or a salt, pro-drug or solvate thereof.

20 In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0;

HET-1 is selected from thiazolyl, thiadiazolyl and pyrazolyl;

25 R^2 is $-\text{CONR}^4\text{R}^5$;

R^4 and R^5 together with the nitrogen to which they are attached form a 6-10 membered bicyclic heterocyclic ring HET-3;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, hydroxymethyl, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

30 or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is a 5- or 6-membered heteroaryl ring;

R^2 is $-S(O)_pR^4$;

5 p is 1 or 2;

R^3 is halo or trifluoromethyl;

R^4 is (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from HET-2, $-OR^5$, $-SO_2R^5$, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R^7) and $-C(O)NR^5R^5$];

10 R^5 is hydrogen or methyl;

HET-2 is a 5- or 6- membered heterocyclyl ring as hereinbefore defined, containing 1 or 2 heteroatoms independently selected from O, N and S; and

R^7 is selected from $-OR^5$ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

15

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0 or 1;

20 HET-1 is a 5- or 6-membered heteroaryl ring;

R^2 is $-S(O)_pR^4$;

p is 1 or 2;

R^3 is halo or trifluoromethyl;

R^4 is (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from

25 HET-2, $-OR^5$, $-SO_2R^5$, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R^7) and $-C(O)NR^5R^5$];

R^5 is hydrogen or methyl;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

30 HET-2 is a 5- or 6- membered heterocyclyl ring as hereinbefore defined, containing 1 or 2 heteroatoms independently selected from O, N and S; and

R^7 is selected from $-OR^5$ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

5 HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

R² is -S(O)_pR⁴;

p is 1 or 2;

R³ is halo or trifluoromethyl;

10 R⁴ is (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from HET-2, -OR⁵, -SO₂R⁵, (3-6C)cycloalkyl and -C(O)NR⁵R⁵];

R⁵ is hydrogen or methyl;

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

15 HET-2 is selected from azetidiny, morpholino, morpholinyl, piperidiny, piperazinyl, 3-oxopiperazinyl, thiomorpholinyl, pyrrolidinyl, pyrrolidonyl, 2,5-dioxopyrrolidinyl, 1,1-dioxotetrahydrothienyl, 2-oxazolidinonyl, 2-oxotetrahydrofuranyl, tetrahydrofuranyl, tetrahydropyranyl, 1,1-dioxothiomorpholino, 1,3-dioxolanyl, 2-oxoimidazolidinyl, 2,4-dioxoimidazolidinyl, pyranyl and 4-pyridonyl; and

20 R⁷ is selected from -OR⁵ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

25 R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

R² is -S(O)_pR⁴;

30 p is 1 or 2;

R³ is halo or trifluoromethyl;

R⁴ is selected from hydrogen, (1-4C)alkyl, [optionally substituted by -OR⁵] and HET-2;

R⁵ is hydrogen or methyl;

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is selected from furyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, pyrrolyl,

5 1,2,4-triazolyl and 1,2,3-triazolyl; and

R⁷ is selected from -OR⁵ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as

10 hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from pyridyl, pyrazinyl, pyridazinyl and pyrimidinyl;

R² is -S(O)pR⁴;

15 p is 1 or 2;

R³ is halo or trifluoromethyl;

R⁴ is (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from HET-2, -OR⁵, -SO₂R⁵, (3-6C)cycloalkyl and -C(O)NR⁵R⁵];

R⁵ is hydrogen or methyl;

20 R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is selected from azetidiny, morpholino, morpholinyl, piperidinyl, piperazinyl, 3-oxopiperazinyl, thiomorpholinyl, pyrrolidinyl, pyrrolidonyl, 2,5-dioxopyrrolidinyl, 1,1-dioxotetrahydrothienyl, 2-oxazolidinonyl, 2-oxotetrahydrofuranlyl, tetrahydrofuranlyl,

25 tetrahydropyranlyl, 1,1-dioxothiomorpholino, 1,3-dioxolanyl, 2-oxoimidazolidinyl, 2,4-dioxoimidazolidinyl, pyranlyl and 4-pyridonyl; and

R⁷ is selected from -OR⁵ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

30 In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from pyridyl, pyrazinyl, pyridazinyl and pyrimidinyl;

R^2 is $-S(O)_pR^4$;

p is 1 or 2;

R^3 is halo or trifluoromethyl;

5 R^4 is selected from hydrogen, (1-4C)alkyl, [optionally substituted by $-OR^5$] and HET-2;

R^5 is hydrogen or methyl;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is selected from furyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl,

10 pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, pyrrolyl, 1,2,4-triazolyl and 1,2,3-triazolyl; and

R^7 is selected from $-OR^5$ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

15 In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl,

20 isoxazolyl and oxadiazolyl;

R^2 is $-S(O)_pR^4$;

p is 1 or 2;

R^3 is halo or trifluoromethyl;

R^4 is (1-4C)alkyl;

25 R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

or a salt, pro-drug or solvate thereof.

30 In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0;

HET-1 is selected from thiazolyl, thiadiazolyl and pyrazolyl;

R^2 is $-S(O)pR^4$;

p is 1 or 2;

R^4 is (1-4C)alkyl;

R^6 is methyl;

5 or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

10 m is 1 and n is 0 or 1;

HET-1 is selected from pyridyl, pyrazinyl, pyridazinyl and pyrimidinyl;

R^2 is $-S(O)pR^4$;

p is 1 or 2;

R^3 is halo or trifluoromethyl;

15 R^4 is (1-4C)alkyl;

R^6 is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

or a salt, pro-drug or solvate thereof.

20 In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R^1 is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is a 5- or 6-membered heteroaryl ring;

25 R^2 is HET-2;

R^3 is halo or trifluoromethyl;

R^5 is hydrogen or (1-4C)alkyl;

HET-2 is a 5- or 6- membered heterocyclyl ring as hereinbefore defined, containing 1 or 2 heteroatoms independently selected from O, N and S; and

30 R^7 is selected from $-OR^5$ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

5 HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

R² is HET-2;

R³ is halo or trifluoromethyl;

R⁵ is hydrogen or methyl;

10 HET-2 is selected from azetidiny, morpholino, morpholinyl, piperidiny, piperazinyl, 3-oxopiperazinyl, thiomorpholinyl, pyrrolidiny, pyrrolidonyl, 2,5-dioxopyrrolidiny, 1,1-dioxotetrahydrothienyl, 2-oxazolidinonyl, 2-oxotetrahydrofuranyl, tetrahydrofuranyl, tetrahydropyranyl, 1,1-dioxothiomorpholino, 1,3-dioxolanyl, 2-oxoimidazolidiny, 2,4-dioxoimidazolidiny, pyranyl and 4-pyridonyl; and

15 R⁷ is selected from -OR⁵ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

20 R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

R² is HET-2;

25 R³ is halo or trifluoromethyl;

R⁵ is hydrogen or methyl;

HET-2 is selected from furyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, pyrrolyl, 1,2,4-triazolyl and 1,2,3-triazolyl; and

30 R⁷ is selected from -OR⁵ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

5 HET-1 is selected from pyridyl, pyrazinyl, pyridazinyl and pyrimidinyl;

R² is HET-2;

R³ is halo or trifluoromethyl;

R⁵ is hydrogen or methyl;

10 HET-2 is selected from azetidiny, morpholino, morpholinyl, piperidiny, piperazinyl, 3-oxopiperazinyl, thiomorpholinyl, pyrrolidinyl, pyrrolidonyl, 2,5-dioxopyrrolidinyl, 1,1-dioxotetrahydrothienyl, 2-oxazolidinonyl, 2-oxotetrahydrofuranly, tetrahydrofuranly, tetrahydropyranly, 1,1-dioxothiomorpholino, 1,3-dioxolany, 2-oxoimidazolidinyl, 2,4-dioxoimidazolidinyl, pyranly and 4-pyridonyl; and

R⁷ is selected from -OR⁵ and (1-4C)alkyl;

15 or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

20 m is 1 and n is 0 or 1;

HET-1 is selected from pyridyl, pyrazinyl, pyridazinyl and pyrimidinyl;

R² is HET-2;

R³ is halo or trifluoromethyl;

R⁵ is hydrogen or methyl;

25 HET-2 is selected from furyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, pyrrolyl, 1,2,4-triazolyl and 1,2,3-triazolyl; and

R⁷ is selected from -OR⁵ and (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

30

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

R² is HET-2;

5 R³ is halo or trifluoromethyl;

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is selected from azetidiny, morpholino, morpholinyl, piperidiny, piperazinyl, 3-oxopiperazinyl, thiomorpholinyl, pyrrolidinyl, pyrrolidonyl, 2,5-dioxopyrrolidinyl, 1,1-

10 dioxotetrahydrothienyl, 2-oxazolidinonyl, 2-oxotetrahydrofuranyl, tetrahydrofuranyl, tetrahydropyranyl, 1,1-dioxothiomorpholino, 1,3-dioxolanyl, 2-oxoimidazolidinyl, 2,4-dioxoimidazolidinyl, pyranyl and 4-pyridonyl; and

R⁷ is (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

15

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

20 HET-1 is selected from thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl and oxadiazolyl;

R² is HET-2;

R³ is halo or trifluoromethyl;

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl,

25 and dimethylaminomethyl;

HET-2 is selected from furyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, pyrrolyl, 1,2,4-triazolyl and 1,2,3-triazolyl; and

R⁷ is (1-4C)alkyl;

30 or a salt, pro-drug or solvate thereof.

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

HET-1 is selected from pyridyl, pyrazinyl, pyridazinyl and pyrimidinyl;

R² is HET-2;

5 R³ is halo or trifluoromethyl;

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

HET-2 is selected from azetidiny, morpholino, morpholinyl, piperidinyl, piperazinyl, 3-oxopiperazinyl, thiomorpholinyl, pyrrolidinyl, pyrrolidonyl, 2,5-dioxopyrrolidinyl, 1,1-

10 dioxotetrahydrothienyl, 2-oxazolidinonyl, 2-oxotetrahydrofuranly, tetrahydrofuranly, tetrahydropyranly, 1,1-dioxothiomorpholino, 1,3-dioxolany, 2-oxoimidazolidinyl, 2,4-dioxoimidazolidinyl, pyranly and 4-pyridonyl; and

R⁷ is (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

15

In a further aspect of the invention is provided a compound of the formula (I) as hereinbefore defined wherein

R¹ is methoxymethyl;

m is 1 and n is 0 or 1;

20 HET-1 is selected from pyridyl, pyrazinyl, pyridazinyl and pyrimidinyl;

R² is HET-2;

R³ is halo or trifluoromethyl;

R⁶ is selected from methyl, ethyl, bromo, chloro, fluoro, aminomethyl, N-methylaminomethyl, and dimethylaminomethyl;

25 HET-2 is selected from furyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, pyrimidinyl, oxazolyl, isoxazolyl, oxadiazolyl, pyrrolyl, 1,2,4-triazolyl and 1,2,3-triazolyl; and

R⁷ is (1-4C)alkyl;

or a salt, pro-drug or solvate thereof.

30

Further preferred compounds of the invention are each of the Examples, each of which provides a further independent aspect of the invention. In further aspects, the present invention also comprises any two or more compounds of the Examples.

In one aspect, particular compounds of the invention comprise any one or more of:

- 3-(4-[(2-methoxyethyl)amino]carbonyl}phenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
- 3-(4-[(1H-imidazol-2-ylmethyl)amino]carbonyl}phenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-(4-[(1-methylpiperidin-4-yl)amino]carbonyl}phenoxy)-N-1,3-thiazol-2-ylbenzamide;
- 3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
- 10 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{4-[(4-methylpiperazin-1-yl)carbonyl]phenoxy}-N-1,3-thiazol-2-ylbenzamide;
- 3-(3-[(2-methoxyethyl)amino]carbonyl}phenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
- 3-(3-[(1H-imidazol-2-ylmethyl)amino]carbonyl}phenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
- 15 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-(3-[(1-methylpiperidin-4-yl)amino]carbonyl}phenoxy)-N-1,3-thiazol-2-ylbenzamide;
- 3-{3-[(dimethylamino)carbonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
- 20 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{4-[(4-methylpiperazin-1-yl)carbonyl]phenoxy}-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[4-(morpholin-4-ylcarbonyl)phenoxy]benzamide;
- 3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 25 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[4-(pyrrolidin-1-ylcarbonyl)phenoxy]benzamide;
- 3-[4-(7-azabicyclo[2.2.1]hept-7-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 30 3-{2-chloro-4-[(dimethylamino)sulfonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[2-chloro-4-(morpholin-4-ylsulfonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

- 3-[(2-chloro-4-[(1-methylethyl)amino]sulfonyl]phenyl]oxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-{[2-chloro-4-([2-(methyloxy)ethyl]amino)sulfonyl]phenyl}oxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 5 3-([2-chloro-4-[(4-methylpiperazin-1-yl)sulfonyl]phenyl]oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-{4-[(dimethylamino)sulfonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-([4-[(4-methylpiperazin-1-yl)sulfonyl]phenyl]oxy)-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 10 3-{4-[(1-methylethyl)amino]sulfonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-(4-{[(2-methoxyethyl)amino]sulfonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 15 3-(4-cyanophenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-{[4-(aminocarbonyl)phenyl]oxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[3-(ethylsulfonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
- 20 3-[4-(ethylsulfonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-{[3-(methylthio)phenyl]oxy}benzamide;
- 25 3-([4-[(1-methylethyl)thio]phenyl]oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[3-(methylsulfonyl)phenoxy]benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[3-(methylsulfinyl)phenoxy]benzamide;
- 30 3-([4-[(1-methylethyl)sulfonyl]phenyl]oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[4-(methylsulfonyl)phenoxy]benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-1,3-thiazol-2-ylbenzamide;
- 5 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(4-methyl-1,3-thiazol-2-yl)benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(5-methyl-1,3-thiazol-2-yl)benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(5-methyl-1,3,4-
- 10 thiadiazol-2-yl)benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide;
- 3-(3,5-difluorophenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide;
- 15 N-(1-ethyl-1H-pyrazol-3-yl)-3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]benzamide;
- 3-(3,5-difluorophenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;
- N-(5-bromopyridin-2-yl)-3-(3,5-difluorophenoxy)-5-[(1S)-2-methoxy-(1-
- 20 methylethyl)oxy]benzamide;
- 3-(3,5-difluorophenoxy)-N-[4-(hydroxymethyl)-1,3-thiazol-2-yl]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-1H-pyrazol-3-ylbenzamide;
- 25 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(5-methyl-1H-pyrazol-3-yl)-5-[4-(methylsulfonyl)phenoxy]benzamide;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-[4-(methoxymethyl)-1,3-thiazol-2-yl]-5-[4-(methylsulfonyl)phenoxy]benzamide;
- 3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(3-methyl-
- 30 1,2,4-thiadiazol-5-yl)benzamide ;
- 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-(4-{[(1-methylpiperidin-4-yl)amino]carbonyl}phenoxy)-N-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide;

3-[4-(azetidin-1-ylcarbonyl)-2-chlorophenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

3-[4-(azetidin-1-ylcarbonyl)-2-fluorophenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

5 3-[4-(azetidin-1-ylcarbonyl)-2-(trifluoromethyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide; and

3-{4-[(dimethylamino)carbonyl]phenoxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide;

or a salt, pro-drug or solvate thereof.

10

The compounds of the invention may be administered in the form of a pro-drug. A pro-drug is a bioprecursor or pharmaceutically acceptable compound being degradable in the body to produce a compound of the invention (such as an ester or amide of a compound of the invention, particularly an in-vivo hydrolysable ester). Various forms of

15 prodrugs are known in the art. For examples of such prodrug derivatives, see:

a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, *et al.* (Academic Press, 1985);

b) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen;

c) H. Bundgaard, Chapter 5 "Design and Application of Prodrugs", by H. Bundgaard

20 p. 113-191 (1991);

d) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);

e) H. Bundgaard, *et al.*, Journal of Pharmaceutical Sciences, 77, 285 (1988); and

f) N. Kakeya, *et al.*, Chem Pharm Bull, 32, 692 (1984).

The contents of the above cited documents are incorporated herein by reference.

25

Examples of pro-drugs are as follows. An in-vivo hydrolysable ester of a compound of the invention containing a carboxy or a hydroxy group is, for example, a pharmaceutically-acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically-acceptable esters for carboxy include

C₁ to C₆alkoxymethyl esters for example methoxymethyl, C₁ to C₆alkanoyloxymethyl esters

30 for example pivaloyloxymethyl, phthalidyl esters,

C₃ to C₈cycloalkoxycarbonyloxy C₁ to C₆alkyl esters for example

1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters, for example 5-methyl-

1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters.

An in-vivo hydrolysable ester of a compound of the invention containing a hydroxy group includes inorganic esters such as phosphate esters (including phosphoramidic cyclic esters) and α -acyloxyalkyl ethers and related compounds which as a result of the in-vivo hydrolysis of the ester breakdown to give the parent hydroxy group/s. Examples of

5 α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxy-methoxy.

A selection of in-vivo hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and N-(dialkylaminoethyl)-N-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl.

10 A suitable pharmaceutically-acceptable salt of a compound of the invention is, for example, an acid-addition salt of a compound of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. In addition a suitable pharmaceutically-acceptable salt of a benzoxazinone derivative of the
15 invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically-acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

20 A further feature of the invention is a pharmaceutical composition comprising a compound of Formula (I) as defined above, or a salt, solvate or prodrug thereof, together with a pharmaceutically-acceptable diluent or carrier.

According to another aspect of the invention there is provided a compound of Formula (I) as defined above for use as a medicament.

25 Further according to the invention there is provided a compound of Formula (I) for use in the preparation of a medicament for treatment of a disease mediated through GLK, in particular type 2 diabetes.

The compound is suitably formulated as a pharmaceutical composition for use in this way.

30 According to another aspect of the present invention there is provided a method of treating GLK mediated diseases, especially diabetes, by administering an effective amount of

a compound of Formula (I) or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.

Specific diseases which may be treated by a compound or composition of the invention include: blood glucose lowering in type 2 Diabetes Mellitus without a serious risk of hypoglycaemia (and potential to treat type 1), dyslipidemia, obesity, insulin resistance, metabolic syndrome X, impaired glucose tolerance.

As discussed above, thus the GLK/GLKRP system can be described as a potential "Diabetesity" target (of benefit in both Diabetes and Obesity). Thus, according to another aspect of the invention there is provided the use of a compound of Formula (I) or salt, solvate or pro-drug thereof, in the preparation of a medicament for use in the combined treatment or prevention of diabetes and obesity.

According to another aspect of the invention there is provided the use of a compound of Formula (I) or salt, solvate or pro-drug thereof, in the preparation of a medicament for use in the treatment or prevention of obesity.

According to a further aspect of the invention there is provided a method for the combined treatment of obesity and diabetes by administering an effective amount of a compound of Formula (I) or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.

According to a further aspect of the invention there is provided a method for the treatment of obesity by administering an effective amount of a compound of Formula (I) or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.

The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

Suitable pharmaceutically acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium carbonate, granulating and disintegrating agents such as corn starch or algenic acid; binding agents such as starch; lubricating agents such as magnesium stearate, stearic acid or talc; 5 preservative agents such as ethyl or propyl p-hydroxybenzoate, and anti-oxidants, such as ascorbic acid. Tablet formulations may be uncoated or coated either to modify their disintegration and the subsequent absorption of the active ingredient within the gastrointestinal tract, or to improve their stability and/or appearance, in either case, using conventional coating agents and procedures well known in the art.

10 Compositions for oral use may be in the form of hard gelatin capsules in which the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions generally contain the active ingredient in finely powdered form 15 together with one or more suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation products of an alkylene oxide with fatty acids (for example polyoxyethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols, for example 20 heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or 25 condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives (such as ethyl or propyl p-hydroxybenzoate, anti-oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening agents (such as sucrose, saccharine or aspartame).

30 Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil (such as arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil (such as liquid paraffin). The oily suspensions may also contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set out above, and flavouring

agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water generally contain the active ingredient together with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients such as sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, or a mineral oil, such as for example liquid paraffin or a mixture of any of these. Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum tragacanth, naturally-occurring phosphatides such as soya bean, lecithin, an esters or partial esters derived from fatty acids and hexitol anhydrides (for example sorbitan monooleate) and condensation products of the said partial esters with ethylene oxide such as polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavouring and preservative agents.

Syrups and elixirs may be formulated with sweetening agents such as glycerol, propylene glycol, sorbitol, aspartame or sucrose, and may also contain a demulcent, preservative, flavouring and/or colouring agent.

The pharmaceutical compositions may also be in the form of a sterile injectable aqueous or oily suspension, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents, which have been mentioned above. A sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in 1,3-butanediol.

Compositions for administration by inhalation may be in the form of a conventional pressurised aerosol arranged to dispense the active ingredient either as an aerosol containing finely divided solid or liquid droplets. Conventional aerosol propellants such as volatile fluorinated hydrocarbons or hydrocarbons may be used and the aerosol device is conveniently arranged to dispense a metered quantity of active ingredient.

For further information on formulation the reader is referred to Chapter 25.2 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition. Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient. For further information on Routes of Administration and Dosage Regimes the reader is referred to Chapter 25.3 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The size of the dose for therapeutic or prophylactic purposes of a compound of the Formula (I) will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well known principles of medicine.

In using a compound of the Formula (I) for therapeutic or prophylactic purposes it will generally be administered so that a daily dose in the range, for example, 0.5 mg to 75 mg per kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous administration, a dose in the range, for example, 0.5 mg to 30 mg per kg body weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.5 mg to 25 mg per kg body weight will be used. Oral administration is however preferred.

The elevation of GLK activity described herein may be applied as a sole therapy or in combination with one or more other substances and/or treatments for the indication being treated. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. Simultaneous treatment may be in a single tablet or in separate tablets. For example in the treatment of diabetes mellitus, chemotherapy may include the following main categories of treatment:

- 1) Insulin and insulin analogues;
- 2) Insulin secretagogues including sulphonylureas (for example glibenclamide, glipizide), prandial glucose regulators (for example repaglinide, nateglinide) ;
- 3) Agents that improve incretin action (for example dipeptidyl peptidase IV inhibitors, and GLP-1 agonists);

- 4) Insulin sensitising agents including PPARgamma agonists (for example pioglitazone and rosiglitazone), and agents with combined PPARalpha and gamma activity;
- 5) Agents that modulate hepatic glucose balance (for example metformin, fructose 1, 6 bisphosphatase inhibitors, glycogen phosphorylase inhibitors, glycogen synthase kinase inhibitors);
- 6) Agents designed to reduce the absorption of glucose from the intestine (for example acarbose);
- 7) Agents that prevent the reabsorption of glucose by the kidney (SGLT inhibitors);
- 8) Agents designed to treat the complications of prolonged hyperglycaemia (for example aldose reductase inhibitors);
- 9) Anti-obesity agents (for example sibutramine and orlistat);
- 10) Anti- dyslipidaemia agents such as, HMG-CoA reductase inhibitors (eg statins); PPAR α agonists (fibrates, eg gemfibrozil); bile acid sequestrants (cholestyramine); cholesterol absorption inhibitors (plant stanols, synthetic inhibitors); bile acid absorption inhibitors (IBATi) and nicotinic acid and analogues (niacin and slow release formulations);
- 11) Antihypertensive agents such as, β blockers (eg atenolol, inderal); ACE inhibitors (eg lisinopril); Calcium antagonists (eg. nifedipine); Angiotensin receptor antagonists (eg candesartan), α antagonists and diuretic agents (eg. furosemide, benzthiazide);
- 12) Haemostasis modulators such as, antithrombotics, activators of fibrinolysis and antiplatelet agents; thrombin antagonists; factor Xa inhibitors; factor VIIa inhibitors; antiplatelet agents (eg. aspirin, clopidogrel); anticoagulants (heparin and Low molecular weight analogues, hirudin) and warfarin;
- 13) Agents which antagonise the actions of glucagon; and
- 14) Anti-inflammatory agents, such as non-steroidal anti-inflammatory drugs (eg. aspirin) and steroidal anti-inflammatory agents (eg. cortisone).

According to another aspect of the present invention there is provided individual compounds produced as end products in the Examples set out below and salts, solvates and pro-drugs thereof.

A compound of the invention, or a salt thereof, may be prepared by any process known to be applicable to the preparation of such compounds or structurally related compounds.

Functional groups may be protected and deprotected using conventional methods. For examples of protecting groups such as amino and carboxylic acid protecting groups (as well as means of formation and eventual deprotection), see T.W. Greene and P.G.M. Wuts,


"Protective Groups in Organic Synthesis", Second Edition, John Wiley & Sons, New York,

5 1991.

Processes for the synthesis of compounds of Formula (I) are provided as a further feature of the invention. Thus, according to a further aspect of the invention there is provided a process for the preparation of a compound of Formula (I), which comprises a process a) to d) (wherein the variables are as defined hereinbefore for compounds of Formula (I) unless


10 otherwise defined):

(a) reaction of an acid of Formula (III) or activated derivative thereof with a compound of Formula (IV),



15 or

(b) reaction of a compound of Formula (V) with a compound of Formula (VI),




wherein X^1 is a leaving group and X^2 is a hydroxyl group or X^1 is a hydroxyl group and


20 X^2 is a leaving group;

process (b) could also be accomplished using the intermediate ester Formula (VII),

wherein P^1 is a protecting group as hereinafter described, followed by ester hydrolysis and amide formation by procedures described elsewhere and well known to those skilled in the art;




(V)




(VII)

or

(c) reaction of a compound of Formula (VIII) with a compound of Formula (IX)



(VIII)




(IX)


wherein X^3 is a leaving group or an organometallic reagent and X^4 is a hydroxyl group or X^3 is a hydroxyl group and X^4 is a leaving group or an organometallic reagent;

process (c) could also be accomplished using the intermediate ester Formula (X),

followed by ester hydrolysis and amide formation by procedures described elsewhere and well known to those skilled in the art;




(VIII)




(X)

or

(d) reaction of a compound of Formula (XI) with a compound of Formula (XII),



(XI)



(XII);

wherein X^5 is a leaving group;

and thereafter, if necessary:

- i) converting a compound of Formula (I) into another compound of Formula (I);
- ii) removing any protecting groups; and/or
- iii) forming a salt, pro-drug or solvate thereof.

Suitable leaving groups X^1 to X^5 for processes b) to d) are any leaving group known in the art for these types of reactions, for example halo, alkoxy, trifluoromethanesulfonyloxy, methanesulfonyloxy, or p-toulenesulfonyloxy; or a group (such as a hydroxy group) that may be converted into a leaving group (such as an oxytriphenylphosphonium group) *in situ*.

Compounds of Formulae (III) to (XII) are commercially available, or are known in the art, or may be made by processes known in the art as shown in the accompanying Examples. For further information on processes for making such compounds, we refer to our PCT publications WO 03/000267, WO 03/015774 and WO 03/000262 and references therein.

Examples of conversions of a compound of Formula (I) into another compound of Formula (I), well known to those skilled in the art, include functional group interconversions such as hydrolysis, oxidation or reduction, and/or further functionalisation by standard reactions such as amide or metal-catalysed coupling, or nucleophilic displacement reactions;

Specific reaction conditions for the above reactions are as follows, wherein when P^1 is a protecting group P^1 is preferably C_{1-4} alkyl, for example methyl or-ethyl:

Process a) – coupling reactions of amino groups with carboxylic acids to form an amide are well known in the art. For example,

- (i) using an appropriate coupling reaction, such as a carbodiimide coupling reaction performed with EDAC in the presence of DMAP in a suitable solvent such as DCM, chloroform or DMF at room temperature; or
- (ii) reaction in which the carboxylic group is activated to an acid chloride by reaction with oxalyl chloride in the presence of a suitable solvent such as methylene chloride. The acid chloride can then be reacted with a compound of Formula (IV) in the presence of a base, such as triethylamine or pyridine, in a suitable solvent such as chloroform or DCM at a temperature between 0°C and room temperature.

Process b) – compounds of Formula (V) and (VI) can be reacted together in a suitable solvent, such as DMF or THF, with a base such as sodium hydride or potassium *tert*-butoxide, at a temperature in the range 0 to 100°C, optionally using metal catalysis such as palladium(II)acetate, palladium on carbon, copper(II)acetate or copper(I)iodide; Alternatively, compounds of Formula (V) and (VI) can be reacted together in a suitable solvent, such as THF

or DCM, with a suitable phosphine such as triphenylphosphine, and azodicarboxylate such as diethylazodicarboxylate;

Process c) - compounds of Formula (VIII) and (IX) can be reacted together in a suitable solvent, such as DMF or THF, with a base such as sodium hydride or potassium *tert*-butoxide, at a temperature in the range 0 to 100°C, optionally using metal catalysis such as palladium(II)acetate, palladium on carbon, copper(II)acetate or copper(I)iodide;

Process d) - reaction of a compound of Formula (XI) with a compound of Formula (XII) can be performed in a polar solvent, such as DMF or a non-polar solvent such as THF with a strong base, such as sodium hydride or potassium *tert*-butoxide at a temperature between 0 and 100°C, optionally using metal catalysis, such as palladium(II)acetate, palladium on carbon, copper(II)acetate or copper(I)iodide.

Certain intermediates of formula (III), (VI), (VII), (IX) and/or (XI) are believed to be novel and comprise an independent aspect of the invention.

Certain intermediates of formula (III), (IX) and/or (XI) wherein R¹ is methoxymethyl are believed to be novel and comprise an independent aspect of the invention.

During the preparation process, it may be advantageous to use a protecting group for a functional group within the molecule. Protecting groups may be removed by any convenient method as described in the literature or known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so as to effect removal of the protecting group with minimum disturbance of groups elsewhere in the molecule.

Specific examples of protecting groups are given below for the sake of convenience, in which "lower" signifies that the group to which it is applied preferably has 1-4 carbon atoms. It will be understood that these examples are not exhaustive. Where specific examples of methods for the removal of protecting groups are given below these are similarly not exhaustive. The use of protecting groups and methods of deprotection not specifically mentioned is of course within the scope of the invention.

A carboxy protecting group may be the residue of an ester-forming aliphatic or araliphatic alcohol or of an ester-forming silanol (the said alcohol or silanol preferably containing 1-20 carbon atoms). Examples of carboxy protecting groups include straight or branched chain (1-12C)alkyl groups (e.g. isopropyl, *t*-butyl); lower alkoxy lower alkyl groups (e.g. methoxymethyl, ethoxymethyl, isobutoxymethyl; lower aliphatic acyloxy lower alkyl groups, (e.g. acetoxymethyl, propionyloxymethyl, butyryloxymethyl, pivaloyloxymethyl);

- lower alkoxycarbonyloxy lower alkyl groups (e.g. 1-methoxycarbonyloxyethyl, 1-ethoxycarbonyloxyethyl); aryl lower alkyl groups (e.g. *p*-methoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, benzhydryl and phthalidyl); tri(lower alkyl)silyl groups (e.g. trimethylsilyl and *t*-butyldimethylsilyl); tri(lower alkyl)silyl lower alkyl groups (e.g. trimethylsilylethyl); and
- 5 (2-6C)alkenyl groups (e.g. allyl and vinyloethyl).

Methods particularly appropriate for the removal of carboxyl protecting groups include for example acid-, metal- or enzymically-catalysed hydrolysis.

- Examples of hydroxy protecting groups include lower alkenyl groups (e.g. allyl); lower alkanoyl groups (e.g. acetyl); lower alkoxycarbonyl groups (e.g. *t*-butoxycarbonyl); lower
- 10 alkenyloxycarbonyl groups (e.g. allyloxycarbonyl); aryl lower alkoxycarbonyl groups (e.g. benzyloxycarbonyl, *p*-methoxybenzyloxycarbonyl, *o*-nitrobenzyloxycarbonyl, *p*-nitrobenzyloxycarbonyl); tri lower alkyl/arylsilyl groups (e.g. trimethylsilyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl); aryl lower alkyl groups (e.g. benzyl) groups; and triaryl lower alkyl groups (e.g. triphenylmethyl).

- 15 Examples of amino protecting groups include formyl, aralkyl groups (e.g. benzyl and substituted benzyl, e.g. *p*-methoxybenzyl, nitrobenzyl and 2,4-dimethoxybenzyl, and triphenylmethyl); di-*p*-anisylmethyl and furylmethyl groups; lower alkoxycarbonyl (e.g. *t*-butoxycarbonyl); lower alkenyloxycarbonyl (e.g. allyloxycarbonyl); aryl lower alkoxycarbonyl groups (e.g. benzyloxycarbonyl, *p*-methoxybenzyloxycarbonyl,
- 20 *o*-nitrobenzyloxycarbonyl, *p*-nitrobenzyloxycarbonyl); trialkylsilyl (e.g. trimethylsilyl and *t*-butyldimethylsilyl); alkylidene (e.g. methylenedene); benzylenedene and substituted benzylenedene groups.

- Methods appropriate for removal of hydroxy and amino protecting groups include, for example, acid-, base, metal- or enzymically-catalysed hydrolysis, or photolytically for groups
- 25 such as *o*-nitrobenzyloxycarbonyl, or with fluoride ions for silyl groups.

- Examples of protecting groups for amide groups include aralkoxymethyl (e.g. benzyloxymethyl and substituted benzyloxymethyl); alkoxymethyl (e.g. methoxymethyl and trimethylsilylethoxymethyl); tri alkyl/arylsilyl (e.g. trimethylsilyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl); tri alkyl/arylsilyloxymethyl (e.g. *t*-butyldimethylsilyloxymethyl,
- 30 *t*-butyldiphenylsilyloxymethyl); 4-alkoxyphenyl (e.g. 4-methoxyphenyl); 2,4-di(alkoxy)phenyl (e.g. 2,4-dimethoxyphenyl); 4-alkoxybenzyl (e.g. 4-methoxybenzyl); 2,4-di(alkoxy)benzyl (e.g. 2,4-di(methoxy)benzyl); and alk-1-enyl (e.g. allyl, but-1-enyl and substituted vinyl e.g. 2-phenylvinyl).

Aralkoxymethyl, groups may be introduced onto the amide group by reacting the latter group with the appropriate aralkoxymethyl chloride, and removed by catalytic hydrogenation. Alkoxymethyl, tri alkyl/arylsilyl and tri alkyl/silyloxymethyl groups may be introduced by reacting the amide with the appropriate chloride and removing with acid; or in the case of the
5 silyl containing groups, fluoride ions. The alkoxyphenyl and alkoxybenzyl groups are conveniently introduced by arylation or alkylation with an appropriate halide and removed by oxidation with ceric ammonium nitrate. Finally alk-1-enyl groups may be introduced by reacting the amide with the appropriate aldehyde and removed with acid.

The following examples are for illustration purposes and are not intended to limit the
10 scope of this application. Each exemplified compound represents a particular and independent aspect of the invention. In the following non-limiting Examples, unless otherwise stated:

(i) evaporations were carried out by rotary evaporation in *vacuo* and work-up procedures were carried out after removal of residual solids such as drying agents by filtration;

15 (ii) operations were carried out at room temperature, that is in the range 18-25°C and under an atmosphere of an inert gas such as argon or nitrogen;

(iii) yields are given for illustration only and are not necessarily the maximum attainable;

(iv) the structures of the end-products of the Formula (I) were confirmed by nuclear
20 (generally proton) magnetic resonance (NMR) and mass spectral techniques; proton magnetic resonance chemical shift values were measured on the delta scale and peak multiplicities are shown as follows: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad; q, quartet, quin, quintet;

(v) intermediates were not generally fully characterised and purity was assessed by
25 thin layer chromatography (TLC), high-performance liquid chromatography (HPLC), infra-red (IR) or NMR analysis; and

(vi) Biotage cartridges refer to pre-packed silica cartridges (from 40g up to 400g), eluted using a biotage pump and fraction collector system; Biotage UK Ltd, Hertford, Herts, UK.


Abbreviations

DCM	dichloromethane;
DEAD	diethylazodicarboxylate;
DIAD	diisopropylazodicarboxylate;
5 DMSO	dimethyl sulphoxide;
DMF	dimethylformamide;
EDAC	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride;
HPLC	high pressure liquid chromatography
10 HPMC	Hydroxypropylmethylcellulose;
LCMS	liquid chromatography / mass spectroscopy;
NMR	nuclear magnetic resonance spectroscopy
RT	room temperature; and
THF	tetrahydrofuran

15

All compound names were derived using ACD NAME computer package.

Example 1: 3-(4-{{(2-Methoxyethyl)amino}carbonyl}phenoxy)-5-(2-(1S)-methoxy-(1-methylethyl)oxy)-N-1,3-thiazol-2-ylbenzamide



20

To a suspension of 4-({3-{{(1S)-2-methoxy-(1-methylethyl)oxy}-5-[(1,3-thiazol-2-ylamino)carbonyl] phenyl}oxy)benzoic acid (107 mg), *O*-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (122 mg) and 2-methoxyethylamine (38 mg) in DMF (2mL), was added diisopropylethylamine (0.11mL) and the mixture stirred at ambient temperature for 1 hour. Water (30mL) was added and the mixture extracted with ethyl acetate (3 x 15mL). The combined organic extracts were washed with brine, dried (MgSO₄), and evaporated to a residue which was chromatographed on silica with ethyl acetate as eluant to give the desired compound (63 mg). ¹H NMR δ (d₆-DMSO): 1.2 (d, 3H), 3.3-(s, 6H), 3.4-3.5

(m, 6H), 4.7-4.8 (m, 1H), 6.85 (s, 1H), 7.1 (d, 2H), 7.25 (m, 2H), 7.55 (d, 2H), 7.9 (d, 2H), 8.45 (s, 1H); m/z 486 (M+H)⁺


In a similar manner to that described above, **Examples 1a-1d** were also prepared:-

Example	Structure	m/z	NMR
1a		508 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.2 (d, 3H), 3.3 (s, 3H), 3.4-3.5 (d, 2H), 4.45 (d, 2H), 4.7-4.8 (m, 1H), 6.85 (m, 3H), 7.1 (d, 2H), 7.25 (d, 2H), 7.55 (d, 2H), 7.95 (d, 2H), 8.95 (t, 1H)
1b		525 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.3 (d, 3H), 1.6-1.75 (m, 4H), 1.95-2.0 (m, 2H), 2.2 (s, 3H), 2.8 (d, 2H), 3.3 (s, 3H), 3.55 (m, 2H), 3.8 (m, 1H), 4.8 (m, 1H), 6.95 (s, 1H), 7.2 (d, 2H), 7.3 (m, 2H), 7.6 (d, 2H), 7.95 (d, 2H), 8.25 (d, 1H)
1c		468 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.25 (d, 3H), 2.2-2.3 (m, 2H), 3.3 (s, 3H), 3.5 (m, 2H), 4.0 (m, 2H), 4.3 (m, 2H), 4.8 (m, 1H), 6.9 (s, 1H), 7.1 (d, 2H), 7.25 (m, 1H), 7.35 (s, 1H), 7.55 (d, 2H), 7.65 (d, 2H), 12.6 (s, 1H)
1d		511 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.25 (d, 3H), 2.2 (s, 3H), 2.3 (m, 4H), 3.3 (s, 3H), 3.4-3.6 (m, 6H), 4.8 (m, 1H), 6.9 (s, 1H), 7.1 (d, 2H), 7.25 (m, 1H), 7.35 (s, 1H), 7.55 (d, 2H), 7.65 (d, 2H), 12.6 (s, 1H)

5


The required acid for Example 1 was prepared as described below:-

4-({3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[(1,3-thiazol-2-ylamino)carbonyl]phenyl}oxy)benzoic acid



A solution of ethyl 4-({3-{[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[(1,3-thiazol-2-ylamino)carbonyl] phenyl}oxy)benzoate (334mg) in THF (10 mL) was added to a solution of lithium hydroxide monohydrate (82mg) in water (5mL). The mixture was stirred at ambient temperature for 16 hours and the THF removed *in vacuo*. The aqueous layer was acidified with 1M hydrochloric acid (1.83mL), and the solid precipitate filtered off, washed with water and dried *in vacuo* to give the desired compound (268mg). ¹H NMR δ (d₆-DMSO): 1.2 (d, 3H), 3.25 (s, 3H), 3.5 (m, 2H), 4.7-4.8 (m, 1H), 6.9 (t, 1H), 7.1 (d, 2H), 7.25 (d, 1H), 7.35 (s, 1H), 7.55 (d, 2H), 7.95 (d, 2H), 12.75 (s, 1H); *m/z* 429 (M+H)⁺


10 Ethyl 4-({3-{[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[(1,3-thiazol-2-ylamino)carbonyl] phenyl}oxy)benzoate



A solution of 3-hydroxy-5-{[(1*S*)-2-methoxy-(1-methylethyl)oxy]-*N*-1,3-thiazol-2-ylbenzamide (1.0g), 4-ethoxycarbonylphenylboronic acid (1.18g), copper (II) acetate (1.19g), triethylamine (2.25mL) and freshly activated 4Å molecular sieves (4g) in dichloromethane (50mL) was stirred at ambient temperature and under ambient atmosphere for 2 days. The reaction mixture was filtered through diatomaceous earth, washed with dichloromethane (2 x 10mL), the dichloromethane removed *in vacuo* and the residual oil partitioned between ethyl acetate (75mL) and 1M hydrochloric acid (30mL). The ethyl acetate layer was separated, washed sequentially with aqueous sodium hydrogen carbonate solution and brine, dried (MgSO₄), and evaporated to a residue which was chromatographed on silica with 30% ethyl acetate in isohexane as eluant to give the desired compound (700mg). ¹H NMR δ (CDCl₃): 1.3 (d, 3H), 1.4 (t, 3H), 3.4 (s, 3H), 3.5-3.6 (m, 2H), 4.35 (q, 2H), 4.5-4.6 (m, 1H), 6.85 (s, 1H), 6.95 (d, 1H), 7.0 (d, 2H), 7.15 (s, 1H), 7.2 (d, 1H), 7.35 (d, 1H), 8.05 (d, 2H); *m/z* 457 (M+H)⁺


25

3-Hydroxy-5-{[(1*S*)-2-methoxy-(1-methylethyl)oxy]-*N*-1,3-thiazol-2-ylbenzamide




A solution of 3-{[(1S)-2-methoxy-(1-methylethyl)oxy]}-5-{[(2-methylphenyl)methyl]oxy}-N-1,3-thiazol-2-ylbenzamide (6.9g) and thioanisole (10mL) in trifluoroacetic acid (65mL) was stirred at ambient temperature for 16 hours. The trifluoroacetic acid was removed *in vacuo* and the residual oil partitioned between ethyl acetate (75mL) and aqueous sodium hydrogen carbonate solution (200mL). The aqueous layer was separated, extracted with ethyl acetate (2 x 75mL), and the combined organic extracts washed with brine, dried (MgSO₄), and evaporated to a residue which was chromatographed on silica with 50% ethyl acetate in isohexane as eluant to give the desired compound (4.6g). ¹H NMR δ (CDCl₃): 1.3 (d, 3H), 3.4 (s, 3H), 3.5-3.6 (m, 2H), 4.5-4.6 (m, 1H), 6.65 (s, 1H), 6.95 (d, 1H), 7.05 (s, 1H), 7.1 (s, 1H), 7.25 (d, 1H); *m/z* 309 (M+H)⁺

3-{[(1S)-2-Methoxy-(1-methylethyl)oxy]}-5-{[(2-methylphenyl)methyl]oxy}-N-1,3-thiazol-2-ylbenzamide



To a solution of 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{[(2-methylphenyl)methyl]oxy} benzoic acid (9.55g) in dichloromethane (140mL) was added oxalyl chloride (2.83mL), followed by DMF (1 drop), and the mixture stirred at ambient temperature for 16 hours. The dichloromethane and excess oxalyl chloride were removed *in vacuo*, the residual oil dissolved in dichloromethane (25mL) and added to a solution of 2-aminothiazole (2.84g) and triethylamine (7.88mL) in dichloromethane (75mL) at 0-5°C, and the mixture stirred at ambient temperature for 4 hours. The dichloromethane and excess triethylamine were removed *in vacuo*, the residual oil partitioned between ethyl acetate (100mL) and 1M hydrochloric acid (100mL). The ethyl acetate layer was separated, washed sequentially with 1M hydrochloric acid, aqueous sodium hydrogen carbonate solution, and brine, dried (MgSO₄), and evaporated to a residue which was chromatographed on alumina with ethyl acetate as eluant to give the desired compound (11.0g). ¹H NMR δ (CDCl₃): 1.3 (d, 3H), 2.35 (s, 3H), 3.4 (s, 3H), 3.5-3.6 (m, 2H), 4.55-4.6 (m, 1H), 5.0 (s, 2H), 6.8 (s, 1H), 6.95 (d, 1H), 7.15 (s, 1H), 7.25 (m, 5H), 7.4 (d, 1H); *m/z* 413 (M+H)⁺


3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[[2-methylphenyl)methyl]oxy}benzoic acid




A solution of methyl 3-[[[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[[2-methylphenyl)methyl]oxy}benzoate (10.65g) in THF (200 mL) and methanol (50mL) was added to a solution of
 5 lithium hydroxide monohydrate (6.0g) in water (100mL). The mixture was stirred at ambient temperature for 16 hours and the THF and methanol removed *in vacuo*. The aqueous layer was acidified to pH1 with hydrochloric acid, and extracted with ethyl acetate (3 x 50mL). The combined organic extracts were washed with brine, dried (MgSO₄), and evaporated to give the desired compound (9.55g); *m/z* 329 (M-H)⁻

10

Methyl 3-[[[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[[2-methylphenyl)methyl]oxy}benzoate




A stirred suspension of methyl 3-hydroxy-5-[[2-methylphenyl)methyl]oxy}benzoate (15.27g) and polymer-supported triphenyl phosphine (39.2g) in dry dichloromethane (900
 15 mL) was cooled in an ice-bath and diisopropyl azodicarboxylate (11.88mL) was added drop wise. The reaction mixture was stirred at 0-5°C for 30 minutes and (*R*)-1-methoxy-propan-2-ol was added dropwise. The reaction mixture was stirred at ambient temperature for 16 hours, filtered through diatomaceous earth and the dichloromethane evaporated to a residue which was chromatographed on silica with 10% ethyl acetate in isohexane as eluant to give the
 20 desired compound (10.65g). ¹H NMR δ (CDCl₃): 1.3 (d, 3H), 2.4 (s, 3H), 3.4 (s, 3H), 3.5-3.6 (m, 2H), 3.9 (s, 3H), 4.55-4.6 (m, 1H), 5.0 (s, 2H), 6.8 (s, 1H), 7.25 (m, 5H), 7.4 (d, 1H)

Methyl 3-hydroxy-5-([(2-methylphenyl)methyl]oxy)benzoate

To a solution of methyl 3,5-dihydroxybenzoate (50g, 0.30M) in DMF (500mL) at 0°C was added sodium hydride (10.8 g, 0.27M) portionwise, maintaining the reaction temperature below 10°C. The reaction was allowed to warm to 15°C, and was stirred for 20 minutes. The mixture was cooled to 0°C and a solution of 2-methylbenzyl bromide (36mL, 0.27M) in DMF (50 mL) was added over 30 minutes. The reaction was warmed to ambient temperature and concentrated *in vacuo*, the residual oil partitioned between ethyl acetate (500mL) and water (250mL), the ethyl acetate layer separated, washed sequentially with water and brine, dried (MgSO₄) and evaporated to a residue which was chromatographed on silica eluting with a gradient of 0-100% ethyl acetate in isohexane to give the desired compound (21.9 g); ¹H NMR δ (CDCl₃) 2.39 (s, 3H), 3.90 (s, 3H), 5.02 (s, 2H), 5.61 (s, 1H), 6.69 (t, 1H), 7.15-7.42 (m, 6H)

15 Example 2: 3-(3-([(2-Methoxyethyl)amino]carbonyl]phenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide



To a suspension of 3-((3-([(1S)-2-methoxy-(1-methylethyl)oxy]-5-[(1,3-thiazol-2-ylamino)carbonyl] phenyl)oxy)benzoic acid (107 mg), *O*-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (122 mg) and 2-methoxyethylamine (38 mg) in DMF (2mL) was added diisopropylethylamine (0.11mL) and the mixture stirred at ambient temperature for 1 hour. Water (30mL) was added and the mixture extracted with ethyl acetate (3 x 15mL). The combined organic extracts were washed with brine, dried (MgSO₄), and evaporated to a residue which was chromatographed on silica, with ethyl acetate as eluant, to give the desired compound (85 mg). ¹H NMR δ (d₆-DMSO): 1.2 (d, 3H), 3.25 (s, 3H), 3.3 (s,

3H), 3.4-3.5 (m, 6H), 4.7-4.8 (m, 1H), 6.8 (s, 1H), 7.2-7.25 (m, 3H), 7.55 (m, 4H), 7.7 (d, 1H) 8.55 (t, 1H), 12.6 (s, 1H); m/z 486 (M+H)⁺


In a similar manner, **Examples 2a-2c** were also prepared:-

5

Example	Structure	m/z	NMR
2a		508 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.25 (d, 3H), 3.3 (s, 3H), 3.5 (m, 2H), 4.45 (d, 2H), 4.7-4.8 (m, 1H), 6.8 (s, 1H), 6.85 (s, 2H), 7.2 (s, 1H), 7.25 (d, 2H), 7.5 (m, 3H), 7.6 (s, 1H), 7.75 (d, 2H), 9.0 (t, 1H)
2b		525 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.25 (d, 3H), 1.5-1.6 (m, 2H), 1.7-1.8 (m, 2H), 1.9-2.0 (m, 2H), 2.15 (s, 3H), 2.75 (d, 2H), 3.3 (s, 3H), 3.5 (m, 2H), 3.6-3.8 (m, 1H), 4.8 (m, 1H), 6.8 (s, 1H), 7.2 (s, 1H), 7.25 (d, 2H), 7.35 (m, 4H), 7.7 (d, 1H), 8.25 (d, 1H)
2c		456 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.2 (d, 3H), 2.9-3.0 (d, 6H), 3.3 (s, 3H), 3.5 (m, 2H), 4.7-4.8 (m, 1H), 6.8 (s, 1H), 7.05 (s, 1H), 7.2 (m, 2H), 7.25 (d, 2H), 7.5 (m, 3H), 12.6 (s, 1H)

The required acid for Example 2 was prepared as described below:-

3-({3-([[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[(1,3-thiazol-2-ylamino)carbonyl]phenyl}oxy)benzoic acid




10

A solution of ethyl 3-({3-([[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[(1,3-thiazol-2-ylamino)carbonyl]phenyl}oxy)benzoate (319mg) in THF (10mL) was added to a solution of lithium hydroxide monohydrate (78mg) in water (5mL). The mixture was stirred at ambient

temperature for 16 hours and the THF removed *in vacuo*. The aqueous layer was acidified with 1M hydrochloric acid (1.75mL), the solid precipitate filtered off, washed with water and dried *in vacuo* to give the desired compound (283mg). ^1H NMR δ (d_6 -DMSO): 1.2 (d, 3H), 3.25 (s, 3H), 3.5 (m, 2H), 4.7-4.8 (m, 1H), 6.85 (t, 1H), 7.25 (m, 2H), 7.35 (dd, 1H), 7.55 (m, 5 4H), 7.75 (d, 1H); m/z 429 ($\text{M}+\text{H}$) $^+$


Ethyl 3-({3-}[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[(1,3-thiazol-2-ylamino)carbonyl]phenyl}oxy)benzoate



- 10 A solution of 3-hydroxy-5-{[(1S)-2-methoxy-(1-methylethyl)oxy]-*N*-1,3-thiazol-2-ylbenzamide (1.0g), 3-ethoxycarbonylphenylboronic acid (1.18g), copper (II) acetate (1.19g), triethylamine (2.25mL) and freshly activated 4Å molecular sieves (4g) in dichloromethane (50mL) was stirred at ambient temperature and under ambient atmosphere for 2 days. The reaction mixture was filtered through diatomaceous earth, washed with dichloromethane (2 x 15 10mL), the dichloromethane removed *in vacuo*, and the residual oil partitioned between ethyl acetate (75mL) and 1M hydrochloric acid (30mL). The ethyl acetate layer was separated, washed sequentially with aqueous sodium hydrogen carbonate solution and brine, dried (MgSO_4), and evaporated to a residue which was chromatographed on silica (eluting with 30% ethyl acetate in isohexane) to give the desired ester (680mg). ^1H NMR δ (CDCl_3): 1.3 20 (d, 3H), 1.4 (t, 3H), 3.4 (s, 3H), 3.5-3.6 (m, 2H), 4.35 (q, 2H), 4.5-4.6 (m, 1H), 6.8 (t, 1H), 6.95 (d, 1H), 7.1 (d, 1H), 7.2 (m, 2H), 7.3 (d, 1H), 7.4 (t, 1H), 7.7 (d, 1H), 7.85 (d, 1H), 11.6 (s, 1H); m/z 457 ($\text{M}+\text{H}$) $^+$

The synthesis of 3-hydroxy-5-{[(1S)-2-methoxy-(1-methylethyl)oxy]-*N*-1,3-thiazol-2-ylbenzamide is described above in Example 1. 25

Example 3: 3-[(1S)-2-Methoxy-(1-methylethyl)oxy]-5-{4-[(4-methylpiperazin-1-yl)carbonyl]phenoxy}-N-(1-methyl-1H-pyrazol-3-yl)benzamide



To a suspension of 4-[(3-{[(1S)-2-methoxy-(1-methylethyl)oxy}-5-{[(1-methyl-1H-pyrazol-3-yl)amino]carbonyl}phenyl)oxy]benzoic acid (212 mg), *O*-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (400 mg) and *N*-Methylpiperazine (105 mg) in DMF (10mL), was added diisopropylethylamine (0.35mL) and the mixture stirred at ambient temperature for 24 hours. Water (30mL) was added and the mixture extracted with ethyl acetate (3 x 15mL). The combined organic extracts were washed with brine, dried (MgSO₄), and evaporated to a residue which was chromatographed on silica eluting with a gradient of 0-50% methanol in ethyl acetate to give the desired compound (130 mg). ¹H NMR δ (CDCl₃): 1.32 (d, 3H), 2.35 (s, 3H), 2.43 (m, 4H), 3.41 (s, 3H), 3.54 (m, 2H), 3.6-3.8 (m, 4H), 3.82 (s, 3H), 4.59 (m, 1H), 6.78 (m, 2H), 7.05 (t, 3H), 7.22 (m, 1H), 7.27 (m, 1H), 7.42 (d, 2H), 8.30 (br s, 1H); *m/z* 508 (M+H)⁺


15

In a similar manner to that described above, **Examples 3a-3d** were also prepared:-

Example	Structure	<i>m/z</i>	NMR
3a		495 (M+H) ⁺	¹ H NMR δ (CDCl ₃): 1.31 (d, 3H), 3.4 (s, 3H), 3.46-3.61 (m, 2H), 3.62-3.77 (m, 8H), 3.81 (s, 3H), 4.60 (m, 1H), 6.78 (m, 2H), 7.02 (s, 1H), 7.07 (m, 2H), 7.22 (m, 1H), 7.28 (m, 1H), 7.42 (d, 2H), 8.31 (br s, 1H)
3b		464 (M+H) ⁺	¹ H NMR δ (CDCl ₃): 1.30 (d, 3H), 2.38 (m, 2H), 3.39 (s, 3H), 3.48-3.60 (m, 2H), 3.78 (s, 3H), 4.20-4.40 (m, 4H), 4.58 (m, 1H), 6.78 (m, 2H), 7.00 (d, 2H), 7.08 (s, 1H), 7.22 (s, 1H), 7.28 (s, 1H), 7.63 (d, 2H), 8.72 (s, 1H)
3c		479 (M+H) ⁺	¹ H NMR δ (CDCl ₃): 1.33 (d, 3H), 1.94 (m, 4H), 3.40 (s, 3H), 3.52 (m, 4H), 3.65 (m, 2H), 3.80 (s, 3H), 4.58 (m, 1H), 6.78 (m, 2H), 7.03 (d, 2H), 7.09 (m, 1H), 7.22 (m, 1H), 7.27 (m, 1H), 7.54 (d, 2H), 8.38 (brs, 1H)
3d		505 (M+H) ⁺	¹ H NMR δ (CDCl ₃): 1.32 (d, 3H), 1.52 (m, 4H), 1.88 (m, 4H), 3.40 (s, 3H), 3.45-3.60 (m, 2H), 3.80 (s, 2H), 4.25 (m, 1H), 4.59 (m, 1H), 4.70 (m, 1H), 6.79 (m, 2H), 7.02 (d, 2H), 7.10 (m, 1H), 7.22 (m, 1H), 7.27 (m, 1H), 7.58 (d, 2H), 8.38 (s, 1H)

The required acid for Example 3 was prepared as described below:-

4-[(3-[(1S)-2-Methoxy-(1-methylethyl)oxy])-5-[(1-methyl-1H-pyrazol-3-yl)amino]carbonyl}phenyl]oxy]benzoic acid




5

A solution of ethyl 4-[(3-[(1S)-2-methoxy-(1-methylethyl)oxy])-5-[(1-methyl-1H-pyrazol-3-yl)amino]carbonyl}phenyl]oxy]benzoate (5.45g) in THF (200 mL) was added to a solution of lithium hydroxide monohydrate (2.52g) in water (100mL). The mixture was stirred at ambient temperature for 48 hours and the THF removed *in vacuo*. The aqueous layer was acidified

10 with 1M hydrochloric acid (60mL), and the solid precipitate filtered off, washed with water and dried *in vacuo* to give the desired acid (5g). ¹H NMR δ (d₆-DMSO): 1.22 (d, 3H), 3.26 (s,


3H), 3.45 (m, 2H), 3.75 (s, 3H), 4.71 (m, 1H), 6.51 (m, 1H), 6.84 (m, 1H), 7.08 (d, 2H), 7.24 (m, 1H), 7.44 (s, 1H), 7.57 (m, 1H), 7.95 (d, 2H), 10.84 (br s, 1H), 12.80 (br s, 1H); m/z 426 (M+H)⁺

5 Ethyl 4-[(3-[(1S)-2-methoxy-(1-methylethyl)oxy])-5-[(1-methyl-1H-pyrazol-3-yl)amino]carbonyl]phenyl]oxy]benzoate



- A solution of 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide (10.0g), 4-ethoxycarbonylphenylboronic acid (9.4g), copper (II) acetate (9g), triethylamine (23mL) and freshly activated 4Å molecular sieves (36g) in dichloromethane (500mL) was stirred at ambient temperature and under ambient atmosphere for 2 days. The reaction mixture was filtered through celite, washed with dichloromethane (2 x 50mL), the dichloromethane removed *in vacuo* and the residual oil partitioned between ethyl acetate (500mL) and 1M hydrochloric acid (200mL). The ethyl acetate layer was separated, washed sequentially with aqueous sodium hydrogen carbonate solution and brine, dried (MgSO₄), and evaporated to a residue which was chromatographed on silica eluting with a gradient of 50-100% ethyl acetate in isohexane to give the desired compound (5.47g). ¹H NMR δ (CDCl₃): 1.3 (m, 3H), 1.41 (t, 3H), 3.39 (s, 3H), 3.49 (m, 1H), 3.58 (m, 1H), 3.78 (s, 3H), 4.38 (q, 2H), 4.58 (m, 1H), 6.79 (m, 2H), 7.01-7.1 (m, 3H), 7.26 (m, 2H), 8.01 (m, 2H), 8.61 (br s, 1H); m/z 454 (M+H)⁺


20 3-Hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide



- To a solution of 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[(phenylmethyl)oxy]benzamide (7.07g) in THF (50mL) and methanol (50mL) was added 10% palladium on carbon (727mg) as a slurry in THF (1 mL) and methanol (1 mL). The mixture was placed under vacuum and stirred under an atmosphere of hydrogen for 70 hours. The mixture was filtered through diatomaceous earth, and the diatomaceous earth washed with

methanol (2 x 100 mL), followed by evaporation *in vacuo*. The residues were dissolved in ethyl acetate (10 mL), treated with isohexane (40 mL), the solid filtered off and washed with isohexane (50 mL) to afford the desired compound (5.17g) which was used without further purification. ^1H NMR δ (d_6 -DMSO): 1.22 (d, 3H), 3.28 (s, 3H, obscured by water), 3.38-3.53 (m, 2H), 3.76 (s, 3H), 4.65 (m, 1H), 6.44 (m, 1H), 6.54 (m, 1H), 6.93 (s, 1H), 7.04 (s, 1H), 7.57 (m, 1H), 9.63 (br s, 1H), 10.60 (s, 1H); m/z 306 ($\text{M}+\text{H}^+$), 304 ($\text{M}-\text{H}^-$)

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[(phenylmethyl)oxy]benzamide




10

A solution of 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[(phenylmethyl)oxy]benzoic acid (8.73g) in dichloromethane (150 mL) was cooled to 0°C . Oxalyl chloride (4.81 mL) and DMF (0.15mL) were slowly added with stirring. The mixture was allowed to warm to ambient temperature and stirred for 16 hours, following which the organics were removed *in vacuo*, and the residues azeotroped with toluene (75mL). The crude material was dissolved in dichloromethane (75 mL) and slowly added to a stirred suspension of 1-methyl-1H-pyrazol-3-amine (3.35g) and diisopropylethylamine (14.4 mL) in dichloromethane (75 mL). The mixture was stirred at ambient temperature for 18 hours, before the organics were evaporated *in vacuo* and the residue dissolved in ethyl acetate (150 mL). The organics were washed with 1M aqueous hydrochloric acid (100 mL) and brine (50 mL), and dried (MgSO_4), before evaporation *in vacuo* to give crude material. This was chromatographed on a 200g Biotage Flash 75 SiO_2 column (eluting with 30 to 90% ethyl acetate in isohexane), and evaporated *in vacuo* to afford the desired compound (7.07g). ^1H NMR δ (d_6 -DMSO): 1.23 (d, 3H), 3.28 (s, 3H, obscured by water), 3.40-3.52 (m, 2H), 3.77 (s, 3H), 4.70 (m, 1H), 5.03 (s, 2H), 6.56 (m, 1H), 6.71 (m, 1H), 7.18 (s, 1H), 7.24 (s, 1H), 7.32-7.47 (br m, 5H), 7.58 (m, 1H), 10.73 (s, 1H); m/z 396 ($\text{M}+\text{H}^+$).


25

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{[phenylmethyl]oxy}benzoic acid




- A solution of methyl 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{[phenylmethyl]oxy} benzoate (77.4 mmol) in a mixture of THF (232 mL) and methanol (232 mL) was treated with a solution of 2M sodium hydroxide (232 mmol), and the reaction mixture stirred for 4 hours at ambient temperature. The resulting solution was diluted with water (250 mL) and most of the organic solvent removed *in vacuo*. The resulting suspension was washed with diethyl ether (3 x 200 mL) and the organic washings discarded. The resulting aqueous solution was acidified to pH4 with 2M hydrochloric acid solution and extracted with ethyl acetate (2 x 200 mL). The extracts were combined, washed with brine, dried (MgSO₄), and evaporated to give the desired compound (99% yield). ¹H NMR δ (d₆-DMSO): 1.20 (d, 3H), 3.46 (m, 2H), 4.64 (m, 1H), 5.15 (s, 2H), 6.83 (app t, 1H), 7.06 (s, 1H), 7.13 (s, 1H), 7.30-7.49 (m, 5H), 12.67 (br s, 1H)

15 Methyl 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{[phenylmethyl]oxy}benzoate



- To a solution of methyl 3-hydroxy-5-{[phenylmethyl]oxy}benzoate (77.4 mmol) in THF was added polymer-supported triphenylphosphine (51.7g of 3 mmol/g loading, 155mmol) and (*R*)-(-)-1-methoxy-2-propanol (102 mmol). The stirred solution was blanketed with argon and cooled in an ice bath. A solution of diisopropylazodicarboxylate (116 mmol) was added dropwise by syringe over 10 minutes. The solution was stirred for 20 minutes and filtered, washing the residue with THF (500 mL). The filtrate and washings were combined, and evaporated to give the desired compound which was used without further purification. ¹H NMR δ (d₆-DMSO): 3.26 (s, 3H), 3.44 (m, 2H), 3.82 (s, 3H), 4.63 (m, 1H), 5.14 (s, 2H), 6.85 (s, 1H), 7.05 (s, 1H), 7.11 (s, 1H), 7.30-7.47 (m, 5H). The ¹H NMR spectrum also contained signals consistent with a small amount of bis(1-methylethyl)hydrazine-1,2-dicarboxylate.

Methyl 3-hydroxy-5-[[phenylmethyl]oxy]benzoate

To a stirred solution of methyl 3,5-dihydroxybenzoate (5.95 mol) in DMF (6 L) was added potassium carbonate (9 mol), and the suspension stirred at ambient temperature under argon.

- 5 To this was added benzyl bromide (8.42 mol) slowly over 1 hour, with a slight exotherm, and the reaction mixture stirred overnight at ambient temperature. The reaction was quenched cautiously with ammonium chloride solution (5 L) followed by water (35 L). The aqueous suspension was extracted with dichloromethane (1 x 3 L and 2 x 5 L). The combined extracts were washed with water (10 L) and dried overnight (MgSO₄). The solution was evaporated in
- 10 *vacuo*, and the crude product chromatographed in 3 batches (flash column, 3 x 2 kg silica, eluting with a gradient consisting of hexane containing 10% dichloromethane, to neat dichloromethane, to dichloromethane containing 50% ethyl acetate) to eliminate starting material. The crude eluant was further chromatographed in 175 g batches (Amicon HPLC, 5 kg normal-phase silica, eluting with isohexane containing 20% v/v of ethyl acetate) to give the
- 15 desired compound (21% yield). ¹H NMR δ (d₆-DMSO): 3.8 (s, 3H), 5.1 (s, 2H), 6.65 (m, 1H), 7.0 (m, 1H), 7.05 (m, 1H), 7.3-7.5 (m, 5H), 9.85 (br s, 1H)

Example 4: General Procedure for Preparation of Halogenated Sulphonamides


- To a solution of the appropriate amine (1.8 mmol) in dichloromethane (2 mL), was added the
- 20 sulphonyl chloride (0.72 mmol) in dichloromethane (2 mL), and the resulting mixture stirred for 18 hours. The mixture was treated with 1M aqueous hydrochloric acid (4 mL) and the organics separated. Evaporation *in vacuo* gave the crude fluorosulphonamide which was used without further purification.

- To a solution of the crude fluorosulphonamide (7.2 mmol) in acetonitrile (3 mL), was added
- 25 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide (0.36 mmol) and potassium carbonate (1.8 mmol). The mixture was heated to 170°C in a 'Smith Creator Microwave' for 100 minutes, before being filtered and the resultant organics evaporated *in vacuo*. The residues were then chromatographed on a Redisep (12g, SiO₂)

cartridge using an Isco Optix chromatography system, eluting with 30 to 100% ethyl acetate in isohexane, and evaporated *in vacuo* to afford the desired compound.

Examples 4a-4e were synthesised using the generic preparation described above:-

Example	Structure	<i>m/z</i>	NMR
4a		523, 525 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.24 (d, 3H), 2.65 (s, 6H), 3.27 (s, 3H, obscured by water), 3.42-3.54 (m, 2H), 3.76 (s, 3H), 4.72-4.81 (m, 1H), 6.55 (m, 1H), 6.93 (m, 1H), 7.20 (d, 1H), 7.26 (s, 1H), 7.48 (s, 1H), 7.58 (m, 1H), 7.70 (dd, 1H), 7.91 (m, 1H), 10.84 (s, 1H)
4b		565, 567 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.24(d, 3H), 2.93 (t, 4H), 3.26 (s, 3H, obscured by water), 3.43-3.54 (m, 2H), 3.63 (t, 4H), 3.77 (s, 3H), 4.78 (m, 1H), 6.54 (m, 1H), 6.95 (m, 1H), 7.22 (d, 1H), 7.28 (s, 1H), 7.50 (m, 1H), 7.58 (m, 1H), 7.68 (dd, 1H), 7.90 (d, 1H), 10.84 (s, 1H)
4c		537, 539 (M+H) ⁺ 535, 537 (M-H) ⁻	¹ H NMR δ (d ₆ -DMSO): 0.95 (d, 6H), 1.23 (d, 3H), 3.27 (s, 3H, obscured by water), 3.27 (m, 1H, obscured by water), 3.42-3.53 (m, 2H), 3.76 (s, 3H), 4.75 (m, 1H), 6.54 (m, 1H), 6.89 (m, 1H), 7.21 (s, 2H), 7.46 (s, 1H), 7.57 (m, 1H), 7.67 (d, 1H), 7.76 (dd, 1H), 7.95 (d, 1H), 10.84 (s, 1H)
4d		553, 555 (M+H) ⁺ 551, 553 (M-H) ⁻	¹ H NMR δ (d ₆ -DMSO): 1.23 (d, 3H), 2.96 (m, 2H), 3.09 (m, 2H), 3.14 (s, 3H), 3.28 (s, 3H, obscured by water), 3.48 (m, 2H), 3.76 (s, 3H), 4.75 (m, 1H), 6.53 (m, 1H), 7.89 (m, 1H), 7.17-7.24 (m, 2H), 7.46 (m, 1H), 7.57 (m, 1H), 7.73 (dd, 1H), 7.82 (m, 1H), 7.96 (m, 1H), 10.84 (s, 1H)

4e ^s		578, 580 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.24 (d, 3H), 2.13 (s, 3H), 2.35 (t, 4H), 2.94 (m, 4H), 3.28 (s, 3H, obscured by water), 3.42-3.53 (m, 2H), 3.75 (s, 3H), 4.76 (m, 1H), 6.54 (m, 1H), 6.93 (m, 1H), 7.21 (d, 1H), 7.28 (s, 1H), 7.49 (s, 1H), 7.58 (m, 1H), 7.69 (dd, 1H), 7.90 (m, 1H), 10.84 (s, 1H)
-----------------	---	-----------------------------	---


^sThe requisite sulphonamide for this example was prepared using a 1:1 ratio of amine : sulphonyl chloride, and isolated by treatment with 1M aqueous sodium hydroxide




- 5 The synthesis of 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide is described in Example 3 above.

Example 5: General Procedure for Preparation of Sulphonamides


- A solution of the requisite chlorosulphonamide from Example 4 above (0.12 mmol) in THF (5 mL) and methanol (5 mL) was treated with 10% palladium on carbon (6 mg) and triethylamine (0.1 mL). The flask was put under vacuum, and stirred under an atmosphere of hydrogen gas. The resulting mixture was stirred at ambient temperature until starting material was consumed, before being filtered through diatomaceous earth and washed with methanol. Evaporation of the organics in vacuo, and azeotroping with diethyl ether (3 x 5mL), followed by drying in vacuo, afforded the desired compound.

Examples 5a-5d were synthesised using the generic procedure described above:-

Example	Structure	<i>m/z</i>	NMR
5a		489 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.23 (d, 3H), 2.60 (s, 6H), 3.27 (s, 3H, obscured by water), 3.43-3.54 (m, 2H), 3.75 (s, 3H), 4.75 (m, 1H), 6.54 (m, 1H), 6.91 (m, 1H), 7.21 (d, 2H), 7.29 (s, 1H), 7.48 (s, 1H), 7.58 (m, 1H), 7.75 (d, 2H), 10.84 (s, 1H)

5b	Chiral 	544 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.24 (d, 3H), 2.13 (s, 3H), 2.36 (s, 4H), 2.88 (s, 4H), 3.28 (s, 3H, obscured by water), 3.43-3.54 (m, 2H), 3.76 (s, 3H), 4.75 (m, 1H), 6.55 (m, 1H), 6.92 (s, 1H), 7.22 (d, 2H), 7.30 (s, 1H), 7.49 (s, 1H), 7.58 (m, 1H), 7.75 (d, 2H), 10.84 (s, 1H)
5c		503 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 0.96 (d, 6H), 1.23 (d, 3H), 3.20 (m, 1H), 3.28 (s, 3H, obscured by water), 3.42-3.53 (m, 2H), 3.76 (s, 3H), 4.55 (m, 1H), 6.55 (m, 1H), 7.88 (m, 1H), 7.18 (d, 2H), 7.25 (s, 1H), 7.46 (s, 1H), 7.50 (d, 1H), 7.59 (m, 1H), 7.81 (d, 2H), 10.84 (s, 1H)
5d		519 (M+H) ⁺ 517 (M-H) ⁻	¹ H NMR δ (d ₆ -DMSO): 1.24 (d, 3H), 2.91 (m, 2H), 3.10 (m, 2H), 3.16 (s, 3H), 3.28 (s, 3H, obscured by water), 3.43-3.52 (m, 2H), 3.76 (s, 3H), 4.76 (m, 1H), 6.53 (m, 1H), 6.87 (m, 1H), 7.19 (d, 2H), 7.25 (s, 1H), 7.45 (s, 1H), 7.57 (m, 1H), 7.64 (m, 1H), 7.80 (d, 2H), 10.84 (s, 1H)

Example 6: 3-(4-Cyanophenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide



- 5 To a stirred solution of 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide (0.164 mmol) in DMF (1 mL) was added a 1M solution of sodium hexamethyldisilazide in THF (0.164 mmol). The reaction was stirred at room temperature for 10 minutes before adding 4-fluorobenzonitrile (0.164 mmol). The reaction was stirred overnight at room temperature, then heated to 60°C and stirred for a further 4 hours. The
- 10 reaction was allowed to cool to room temperature, and treated with a further 0.2 equivalents of 4-fluorobenzonitrile and sodium hexamethyldisilazide, heated to 70°C and stirred at this temperature for 3 hours. The reaction was cooled to room temperature, and treated with a further 0.2 equivalents of sodium hexamethyldisilazide, warmed to 70°C, and stirred at this temperature overnight. The solvent was removed *in vacuo* and the residual oil partitioned


between ethyl acetate and water. The water layer was separated and re-extracted with ethyl acetate. The combined organic layers were washed with brine, dried (MgSO_4), filtered and evaporated to a residue which was chromatographed on silica, using 0-1% methanol in dichloromethane as the eluent, to give the desired product (60% yield). ^1H NMR δ (CDCl_3):

5 1.35 (d, 3H), 3.40 (s, 3H), 3.55 (m, 2H), 3.78 (s, 3H), 4.60 (m, 1H), 6.80 (m, 2H), 7.10 (m, 3H), 7.30 (m, 2H), 7.62 (d, 2H), 8.55 (br s, 1H); m/z 407 ($\text{M}+\text{H}^+$), 405 ($\text{M}-\text{H}^-$)

The synthesis of 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide is described in Example 3 above.


10

Example 7: 3-{[4-(Aminocarbonyl)phenyl]oxy}-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide




A suspension of 3-(4-cyanophenoxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide (0.25 mmol), prepared as described in Example 6 above, sodium azide (0.28 mmol) and zinc bromide (0.25 mmol) in water (2mL) was heated to reflux and stirred at this temperature overnight. Isopropanol (2 mL) was added, and the reaction heated at reflux for a further 24 hours. The reaction was cooled to room temperature, evaporated to half volume *in vacuo*, and the residue partitioned between ethyl acetate and water. The water layer was separated and re-extracted with ethyl acetate. The combined organic layers were washed with brine, dried (MgSO_4), filtered and evaporated to a residue which was chromatographed on silica with 0-10% methanol in dichloromethane as eluent to yield crude material. This material was dissolved in ethyl acetate and washed twice with 2M sodium hydroxide. The organic layer was washed with brine, dried (MgSO_4), filtered and evaporated. This material was dissolved in dichloromethane and purified using an 'Isolute-NH2' ion-exchange column eluting with 10% methanol:dichloromethane to yield the desired product. ^1H NMR δ (CDCl_3): 1.30 (d, 3H), 3.40 (s, 3H), 3.50 (m, 2H), 3.75 (s, 3H), 4.60 (m, 1H), 6.80 (m, 2H), 7.00 (d, 2H), 7.05 (s, 1H), 7.25 (m, 2H), 7.80 (d, 2H), 8.75 (br s, 1H); m/z 423 ($\text{M}-\text{H}^-$)

Example 8: 3-[3-(Ethylsulfonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide



A solution of 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide (154 mg), 3-ethanesulphonylbenzeneboronic acid (203 mg), copper(II) acetate (183 mg), triethylamine (0.345mL) and freshly activated 4Å molecular sieves (1g) in dichloromethane (10mL), was stirred at ambient temperature and under ambient atmosphere for 3 days. The reaction mixture was filtered through diatomaceous earth, washed with dichloromethane (2 x 10mL), the dichloromethane removed *in vacuo* and the residual oil partitioned between ethyl acetate (50mL) and 1M hydrochloric acid (35mL). The ethyl acetate layer was separated, washed sequentially with saturated aqueous sodium hydrogen carbonate and brine, dried (MgSO₄) and evaporated to a residue which was chromatographed on alumina with 5% methanol in ethyl acetate as eluant. Further chromatography on silica with 50% ethyl acetate in isohexane as eluant gave the desired compound (108 mg). ¹H NMR δ (CDCl₃): 1.2-1.35 (m, 6H), 3.15 (q, 2H), 3.4 (s, 3H), 3.5-3.6 (m, 2H), 4.5-4.6 (m, 1H), 6.8 (s, 1H), 7.0 (d, 1H), 7.2 (d, 1H), 7.25 (d, 1H), 7.3 (d, 1H), 7.35 (d, 1H), 7.55 (d, 2H), 8.05 (d, 1H); *m/z* 477 (M+H)⁺

The following compounds were also prepared in an analogous fashion from 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide, or from 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide:

Example	Structure	<i>m/z</i>	NMR
8a		477 (M+H) ⁺	¹ H NMR δ (CDCl ₃): 1.2-1.35 (m, 6H), 3.15 (q, 2H), 3.4 (s, 3H), 3.5-3.6 (m, 2H), 4.5-4.6 (m, 1H), 6.8 (s, 1H), 6.95 (d, 1H), 7.2 (d, 2H), 7.25 (d, 2H), 7.4 (s, 1H), 7.85 (d, 2H)


8b		428 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.22 (d, 3H), 2.45 (s, 3H), 3.30 (s, 3H), 3.46 (m, 2H), 3.76 (s, 3H), 4.72 (m, 1H), 6.53 (m, 1H), 6.72 (m, 1H), 6.80 (m, 1H), 6.94 (m, 1H), 7.05 (d, 1H), 7.12 (s, 1H), 7.34 (t, 1H), 7.39 (s, 1H), 7.57 (m, 1H), 10.81 (bs, 1H)
8c		456 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.11 (m, 9H), 3.26 (s, 3H), 3.37 (m, 1H), 3.45 (m, 2H), 3.76 (s, 3H), 4.72 (m, 1H), 6.53 (m, 1H), 6.74 (m, 1H), 7.02 (d, 2H), 7.16 (s, 1H), 7.39 (m, 1H), 7.42 (d, 2H), 7.57 (m, 1H), 10.81 (bs, 1H)

The syntheses of 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-1,3-thiazol-2-ylbenzamide and 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide are described in Examples 1 and 3 respectively.

5

Example 9a: 3-[(1S)-2-Methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[3-(methylsulfonyl)phenoxy]benzamide

Example 9b: 3-[(1S)-2-Methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[3-(methylsulfinyl)phenoxy]benzamide



10


To a solution of 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)-5-[3-(methylthio)phenyl]oxy}benzamide (prepared as described in Example 8b above, 270mg) in dichloromethane (5 mL) was added *m*-chloroperbenzoic acid (1.3 equivalents) and the reaction stirred at room temperature for 1 hour. A further 1.4 equivalents of *m*-

15 chloroperbenzoic acid was added, and the reaction stirred at room temperature for a further 30 minutes. The reaction was added to saturated aqueous sodium metabisulphite and stirred for 20 minutes. The organic layer was separated, washed with brine, dried (MgSO₄), and evaporated to a white foam. The crude mixture was purified using a 20g Redisep column eluting with 0-5% methanol in dichloromethane to yield the desired sulphone (117mg). ¹H

NMR δ (d_6 -DMSO): 1.12 (d, 3H), 3.22 (s, 3H), 3.26 (s, 3H), 3.47 (m, 2H), 3.75 (s, 3H), 4.75 (m, 1H), 6.54 (m, 1H), 6.85 (m, 1H), 7.23 (s, 1H), 7.40 (m, 1H), 7.45 (s, 1H), 7.52 (m, 1H), 7.57 (m, 1H), 7.68 (m, 2H), 10.84 (br s, 1H); m/z 460 ($M+H$)⁺

A further fraction yielded the desired sulfoxide (105mg). ¹H NMR δ (d_6 -DMSO): 1.12 (d, 3H), 2.75 (s, 3H), 3.26 (s, 3H), 3.47 (m, 2H), 3.76 (s, 3H), 4.73 (m, 1H), 6.53 (m, 1H), 6.80 (m, 1H), 7.19 (m, 2H), 7.33 (m, 1H), 7.44 (m, 2H), 7.59 (m, 2H), 10.83 (br s, 1H); m/z 444 ($M+H$)⁺

Example 10: 3-({4-[(1-methylethyl)sulfonyl] phenyl}oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide



In a similar manner to that described above for Example 9, 3-({4-[(1-methylethyl)sulfonyl] phenyl}oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide was prepared from 3-({4-[(1-methylethyl)thio]phenyl}oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide. ¹H NMR δ (d_6 -DMSO): 1.32 (m, 9H), 3.27 (m, 1H), 3.41 (s, 3H), 3.50 (dd, 1H), 3.58 (dd, 1H), 3.80 (s, 3H), 4.61 (m, 1H), 6.82 (m, 2H), 7.09 (d, 2H), 7.17 (m, 1H), 7.28 (m, 1H), 7.33 (m, 1H), 7.84 (d, 2H), 8.86 (br s, 1H); m/z 488 ($M+H$)⁺






The synthesis of 3-({4-[(1-methylethyl)thio]phenyl}oxy)-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide is described in Example 8c above.

Example 11: General Procedure for Amide Synthesis – HATU Coupling

Diisopropylethylamine (2.5 equivalents) was added to a suspension of 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{{4-(methylsulfonyl)phenyl}oxy}benzoic acid (1 equivalent), *O*-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (1.25-equivalents) and amine (1.25 equivalents) in DMF (20mL). The initial suspension dissolved into a dark orange solution. The resulting mixture was stirred at ambient temperature for 2 hours. The DMF was removed *in vacuo*, and the residue azeotroped with toluene. Water was added and

the mixture extracted with ethyl acetate. The extracts were combined and washed sequentially with 1M hydrochloric acid, saturated sodium hydrogen carbonate solution and brine. The solution was dried (MgSO_4), filtered, and evaporated *in vacuo* to give the crude product which was chromatographed (50% ethyl acetate in isohexane) to give desired compound (40-70% 5 yield).

Examples 11a-11h were prepared using an analogous method to that described above from the appropriate acid and amino heterocycle:

Example	Structure	<i>m/z</i>	NMR
11a		460 ($\text{M}+\text{H}^+$)	^1H NMR δ (d_6 -DMSO): 1.2 (d, 3H), 3.2 (s, 3H), 3.25 (s, 3H), 3.5 (m, 2H), 3.8 (s, 3H), 4.75 (m, 1H), 6.55 (s, 1H), 6.9 (s, 1H), 7.2 (d, 2H), 7.3 (s, 1H), 7.45 (s, 1H), 7.6 (s, 1H), 7.9 (d, 2H), 10.85 (br s, 1H)
11b		463 ($\text{M}+\text{H}^+$) 461 ($\text{M}-\text{H}^-$)	^1H NMR δ (d_6 -DMSO): 1.2 (d, 3H), 3.2 (s, 3H), 3.25 (s, 3H), 3.5 (m, 2H), 4.75 (m, 1H), 6.9 (s, 1H), 7.2 (d, 2H), 7.3 (s, 1H), 7.4 (s, 1H), 7.55 (d, 1H), 7.6 (s, 1H), 7.9 (d, 2H), 12.6 (br s, 1H)
11c		477 ($\text{M}+\text{H}^+$) 475 ($\text{M}-\text{H}^-$)	^1H NMR δ (d_6 -DMSO): 1.2 (d, 3H), 2.25 (s, 3H), 3.2 (s, 3H), 3.25 (s, 3H), 3.5 (m, 2H), 4.75 (m, 1H), 6.8 (s, 1H), 6.95 (s, 1H), 7.2 (d, 2H), 7.3 (s, 1H), 7.4 (s, 1H), 7.95 (d, 2H), 12.6 (br s, 1H)
11d		477 ($\text{M}+\text{H}^+$) 475 ($\text{M}-\text{H}^-$)	^1H NMR δ (d_6 -DMSO): 1.2 (d, 3H), 2.4 (s, 3H), 3.2 (s, 3H), 3.25 (s, 3H), 3.5 (m, 2H), 4.75 (m, 1H), 6.95 (s, 1H), 7.2 (s, 1H), 7.25 (d, 2H), 7.4 (s, 1H), 7.6 (s, 1H), 7.95 (d, 2H), 12.4 (br s, 1H)
11e		478 ($\text{M}+\text{H}^+$) 476 ($\text{M}-\text{H}^-$)	^1H NMR δ (d_6 -DMSO): 1.2 (d, 3H), 2.6 (s, 3H), 3.2 (s, 3H), 3.25 (s, 3H), 3.5 (m, 2H), 4.75 (m, 1H), 7.0 (s, 1H), 7.2 (d, 2H), 7.4 (s, 1H), 7.6 (s, 1H), 7.95 (d, 2H)

11f*		478 (M+H) ⁺ 476 (M-H) ⁻	¹ H NMR δ (d ₆ -DMSO): 1.2 (d, 3H), 2.5 (s, 3H), 3.2 (s, 3H), 3.25 (s, 3H), 3.5 (m, 2H), 4.75 (m, 1H), 7.0 (s, 1H), 7.2 (d, 2H), 7.4 (s, 1H), 7.6 (s, 1H), 7.95 (d, 2H), 13.5 (br s, 1H)
11g^{\$}		474 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.24 (d, 3H), 1.38 (t, 3H), 3.20 (s, 3H), 3.30 (s, 3H), 3.51 (m, 2H), 4.06 (s, 3H), 4.79 (m, 1H), 6.58 (s, 1H), 6.92 (s, 1H), 7.26 (d, 2H), 7.30 (s, 1H), 7.50 (s, 1H), 7.56 (s, 1H), 7.96 (d, 2H), 10.89 (s, 1H)
11h		421 (M+H) ⁺ 419 (M-H) ⁻	¹ H NMR δ (d ₆ -DMSO): 1.24 (d, 3H), 3.29 (s, 3H obscured by solvent peak), 3.48 (m, 2H), 4.76 (m, 1H), 6.83 (dd, 2H), 6.94 (m, 1H), 7.04 (m, 1H), 7.27 (m, 1H), 7.33 (m, 1H), 7.54 (d, 2H), 12.62 (bs, 1H)

^{\$}The required amino pyrazole was prepared as follows. Sodium hydride (60% dispersion in mineral oil, 39 mg, 0.973 mmol), was added to 5-nitro-1*H*-pyrazole (100 mg, 0.885 mmol) in dry DMF (2 mL) under an argon atmosphere. The solution was stirred for 5 minutes, then ethyl iodide (0.85 mL, 1.062 mmol) added and the reaction warmed to 80°C for 3 hours.

- 5 Saturated aqueous sodium hydrogen carbonate (30mL) was added, and the mixture extracted with diethyl ether (40 mL). The combined organic extracts were washed with brine, dried (MgSO₄), and evaporated to a residue which was purified by chromatography on silica (eluting with isohexane containing ethyl acetate, 33% v/v) to give the alkylated pyrazole (80 mg) which was used in the next step without further purification. ¹H NMR δ (CDCl₃): 1.58 (t, 10 3H), 4.26 (q, 2H), 6.91 (d, 1H), 7.48 (d, 1H).


To a solution of the alkylated pyrazole (70 mg, 0.50 mmol) in THF (5 mL) under an inert atmosphere was added 10% palladium on carbon (15 mg). The flask was evacuated and refilled 3 times with hydrogen gas, and stirred vigorously at room temperature for 3 hours. The reaction mixture was refilled with argon, and a further portion of 10% palladium on 15 carbon (50 mg) added, followed by refilling as above with a hydrogen atmosphere. The reaction was stirred for 16 hours, filtered through diatomaceous earth, and evaporated to afford the title compound (56 mg) as a colourless oil which was used without further

purification. ^1H NMR δ (CDCl_3): 1.42 (t, 3H), 3.58 (br. s, 2H), 3.98 (q, 2H), 5.59 (d, 1H), 7.16 (d, 1H)

*Example 11f may be crystallised by allowing isohexane to vapour diffuse into a solution of the compound in ethylacetate, in a closed system, with subsequent slow evaporation of the mixture at room temperature over 4 days, mp 109-112.


The required acids for Examples 11a-11h were prepared as described below:-

3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenyl]oxy} benzoic acid



- 10 A solution of methyl 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenyl]oxy} benzoate (60.9 mmol) in THF (400 mL) was treated with a solution of 1M sodium hydroxide (125 mmol), and the reaction mixture stirred for 13 hours at ambient temperature. Most of the organic solvent was removed *in vacuo*, and the remaining solution was diluted with water (150 mL). The resulting aqueous solution was acidified to pH4 with 1M citric acid solution, and extracted with ethyl acetate (2 x 100 mL). The extracts were combined, washed with brine, dried (MgSO_4), and evaporated to give the desired compound (83% yield). ^1H NMR δ (d_6 -DMSO): 1.2 (d, 3H), 3.2 (s, 3H), 3.26 (s, 3H), 3.44 (m, 2H), 4.63 (m, 1H), 7.05 (s, 1H), 7.11 (s, 1H), 7.2 (d, 2H), 7.3 (s, 1H), 7.9 (d, 2H); m/z 479 (M-H)⁺

Methyl 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenyl]oxy} benzoate



20


- A suspension of methyl 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]benzoate (154 mmol), boronic acid (1.1 equivalents), copper (II) acetate (1.1 equivalents), triethylamine (5 equivalents) and freshly activated 4Å molecular sieves (200 g) in dichloromethane (500 mL) was stirred at ambient temperature and under ambient atmosphere for 2 days. The reaction mixture was filtered, the dichloromethane removed *in vacuo* and the residual oil partitioned between ethyl acetate and 1-2M hydrochloric acid. The ethyl acetate layer was separated, washed with aqueous sodium hydrogen carbonate and brine, dried (MgSO_4), and evaporated

25

to a residue which was chromatographed on silica (with 20-60% ethyl acetate in isohexane as eluant) to give the desired ester (58% yield). $^1\text{H NMR } \delta$ (d_6 -DMSO): 1.2 (d, 3H), 3.2 (s, 3H), 3.26 (s, 3H), 3.44 (m, 2H), 3.8 (s, 3H), 4.65 (m, 1H), 7.05 (s, 1H), 7.11 (s, 1H), 7.2 (d, 2H), 7.3 (s, 1H), 7.9 (d, 2H)

5


Methyl 3-Hydroxy-5-[(1*S*)-2-methoxy-(1-methylethyl)oxy]benzoate



Methyl 3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[[phenylmethyl]oxy]benzoate (50.0g; 0.152mmol) was dissolved in a mixture of THF:ethanol (600mL) and the flask evacuated and 10 purged with nitrogen (3 times). 10% Palladium on carbon (5.0g) was added and the flask further evacuated and finally purged with hydrogen gas. The reaction mixture was stirred at ambient temperature for 20 hours until completion. The reaction mixture was evacuated and purged with nitrogen (3 times). The catalyst was filtered off, and the filtrate concentrated *in vacuo* to give the desired compound (36.7g). $^1\text{H NMR } \delta$ (d_6 -DMSO): 1.2 (d, 3H), 3.25 (s, 15 3H), 3.44 (m, 2H), 3.82 (s, 3H), 4.55 (m, 1H), 6.6 (s, 1H), 6.9 (s, 1H), 6.95 (s, 1H), 9.8 (s, 1H)

The synthesis of methyl 3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[[phenylmethyl]oxy]benzoate is described above in Example 1.


3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-{3,5-difluorophenoxy}benzoic acid




20

This was prepared from methyl 3-[(3,5-difluorophenyl)oxy]-5-[(1*S*)-2-methoxy-(1-methylethyl)oxy]benzoate using an analogous procedure to that described above for the synthesis of 3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[[4-(methylsulfonyl)phenyl]oxy]benzoic acid:-

25 $^1\text{H NMR } \delta$ (d_6 -DMSO): 1.21 (d, 3H), 3.26 (s, 3H obscured by solvent peak), 3.46 (m, 2H), 4.67 (m, 1H), 6.81 (d, 2H), 6.96 – 7.08 (m, 3H), 7.27 (s, 1H), 13.13 (bs, 1H); m/z 337 (M-H)⁻


Methyl 3-[(3,5-difluorophenyl)oxy]-5-[(1*S*)-2-methoxy-(1-methylethyl)oxy]benzoate

To a solution of methyl 3-[(3,5-difluorophenyl)oxy]-5-hydroxybenzoate (15.0 mmol), (*R*)-(-)-1-methoxy-2-propanol (18.75 mmol) and triphenylphosphine (18.0 mmol) in anhydrous THF 5 (100 mL) at 0°C was added diisopropylazodicarboxylate (18.0 mmol). The reaction was stirred at ambient temperature overnight, concentrated *in vacuo*, and the residue triturated with a 1:1 mixture of ethyl acetate:isohexane. The solid was removed by filtration and the filtrate concentrated *in vacuo*, chromatographed on silica (using a Biotage Flash 75 eluting with 10-15% ethyl acetate in isohexane) to give the title compound (75% yield). ¹H NMR δ (d₆-10 DMSO): 1.21 (d, 3H), 3.27 (s, 3H obscured by solvent peak), 3.46 (m, 2H), 3.82 (s, 3H), 4.69 (m, 1H), 6.81 (dd, 2H), 7.01-7.07 (m, 2H), 7.10 (s, 1H), 7.28 (s, 1H)

Methyl 3-[(3,5-difluorophenyl)oxy]-5-hydroxybenzoate


15 To a solution of methyl 3-[(3,5-difluorophenyl)oxy]-5-[[4-methylphenyl)sulfonyl]oxy]benzoate (16.3 mmol) in methanol (60 mL) was added a 20% solution of potassium hydroxide in methanol (13.75 g). The mixture was heated at 50°C for 1 hour then allowed to cool. Water (20 mL) was added and the mixture immediately acidified with 1M hydrochloric acid. The methanol was removed *in vacuo* and the residue extracted with ethyl acetate. The organic 20 phase was separated, washed with brine, dried (MgSO₄), and concentrated *in vacuo* to give the title compound (92% yield). ¹H NMR δ (d₆-DMSO): 3.80 (s, 3H), 6.72 (m, 1H), 6.79 (m, 2H), 6.98 – 7.05 (m, 2H), 7.19 (m, 1H), 10.18 (bs, 1H); *m/z* 279 (M-H)⁺

Methyl 3-[(3,5-difluorophenyl)oxy]-5-[[[(4-methylphenyl)sulfonyl]oxy]benzoate



To a solution of methyl 3-hydroxy-5-[[[(4-methylphenyl)sulfonyl]oxy]benzoate (30 mmol), copper (II) acetate (36 mmol), 3,5-difluorophenylboronic acid (42 mmol) and 4Å molecular
 5 sieves (30 g) in dichloromethane (300 mL) was added triethylamine (150 mmol). The reaction was allowed to stir for 40 hours then filtered and concentrated *in vacuo*. The residue was dissolved in ethyl acetate, washed with 1M citric acid solution, 1M sodium hydrogen carbonate solution and brine, then dried (MgSO₄) and concentrated *in vacuo*. The residue was chromatographed on silica (Biotage Flash 75) eluting with 10- 25% ethyl acetate in isohexane
 10 to give the title compound (55% yield). ¹H NMR δ (d₆-DMSO): 2.39 (s, 3H), 3.83 (s, 3H), 6.74 (dd, 2H), 6.93 (m, 1H), 7.08 (m, 1H), 7.44 (m, 3H), 7.50 (s, 1H), 7.74 (d, 2H); *m/z* 452 (M+NH₄)⁺, 433 (M-H)⁻

Methyl 3-hydroxy-5-[[[(4-methylphenyl)sulfonyl]oxy]benzoate





15 Methyl 3,5-dihydroxybenzoate (0.40g) and 4-toluenesulphonylchloride (0.45g) was stirred vigorously in diethyl ether (20mL) with saturated aqueous sodium hydrogen carbonate (20mL) at ambient temperature for 62 hours. The aqueous layer was removed and the residue washed sequentially with saturated aqueous sodium hydrogen carbonate, brine, dried (MgSO₄),
 20 filtered, and concentrated *in vacuo* to yield a colourless oil. The crude product was dissolved in diethyl ether, washed with saturated aqueous potassium carbonate then with brine, dried (MgSO₄), filtered and concentrated *in vacuo* to give a colourless oil which crystallised on standing to give the title compound (0.51g). ¹H NMR δ (d₆-DMSO): 2.43 (s, 3H), 3.82 (s, 3H), 6.66 (m, 1H), 6.97 (s, 1H), 7.26 (s, 1H), 7.47 (d, 2H), 7.75 (d, 2H); *m/z* 340 (M+NH₄)⁺

Example 12: General Procedure for Amide Synthesis – Oxalyl Chloride Coupling

To a stirred solution of 3-{(1S)-2-methoxy-(1-methylethyl)oxy}-5-{3,5-difluorophenoxy} benzoic acid (0.285mmol) in dry dichloromethane (2mL), was added, dropwise under argon, oxalyl chloride (2 equivalents) and DMF (1 drop). The resulting solution was stirred at ambient temperature for 1-2 hrs. The solvent was removed *in vacuo* and the crude mixture taken up in pyridine (2mL) and added to the appropriate amine (2.2 equivalents). The reaction mixture was stirred at room temperature, or heated if necessary, and monitored by TLC and/or LCMS. The pyridine was removed *in vacuo*, and water and ethyl acetate added. The organic layer was washed sequentially with 1M citric acid and brine solution and dried (MgSO₄), concentrated *in vacuo*, and the residue chromatographed on silica (eluting with 30-90% ethyl acetate in isohexane) to give the desired product (typically 35-40% yield).


Examples 12a & 12b were prepared using the appropriate amine:

12a		419 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.23 (d, 3H), 3.27 (s, 3H obscured by solvent peak), 3.47 (m, 2H), 3.76 (s, 3H), 4.74 (m, 1H), 6.55 (d, 1H), 6.80 (d, 2H), 6.86 (m, 1H), 7.02 (m, 1H), 7.24 (s, 1H), 7.44 (s, 1H), 7.57 (s, 1H), 10.82 (br s, 1H)
12b^s		493, 495 (M+H) ⁺	¹ H NMR δ (d ₆ -DMSO): 1.23 (d, 3H), 3.26 (s, 3H), 3.47 (m, 2H), 4.76 (m, 1H), 6.80 (dd, 2H), 6.92 (t, 1H), 7.02 (m, 1H), 7.26 (m, 1H), 7.45 (m, 1H), 8.04 (m, 1H), 8.13 (d, 1H), 8.49 (m, 1H), 11.01 (br s, 1H)

^sIn this example, the acid chloride was taken up in THF, followed by addition of pyridine and the appropriate amine.


The synthesis of 3-{(1S)-2-methoxy-(1-methylethyl)oxy}-5-{3,5-difluorophenoxy}benzoic acid is described in Example 11 above.

Example 13: 3-(3,5-Difluorophenoxy)-N-[4-(hydroxymethyl)-1,3-thiazol-2-yl]-5-[2-(1S)-methoxy-(1-methylethyl)oxy]benzamide



To a solution of 3-(3,5-difluorophenoxy)-N-[4-chloromethyl-1,3-thiazol-2-yl]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]benzamide (0.107 mmol) in THF (1 mL) was added 0.5M sodium hydroxide solution (1 mL). The reaction was stirred at ambient temperature for 2 hours and the organics removed *in vacuo*. The residue was acidified with 1M citric acid and partitioned between ethyl acetate and water. The organic phase was separated, dried (MgSO₄), and concentrated *in vacuo*. The residue was chromatographed on silica eluting with 80% ethyl acetate in isohexane to give the title compound which was precipitated from a concentrated diethyl ether solution by the addition of isohexane to give a solid sample (35% yield). ¹H NMR δ (d₆-DMSO): 1.24 (d, 3H), 3.28 (s, 3H obscured by solvent peak), 3.48 (m, 2H), 4.49 (s, 2H), 4.75 (m, 1H), 6.83 (d, 2H), 6.93 (s, 1H), 6.98 (s, 1H), 7.04 (m, 1H), 7.32 (s, 1H), 7.54 (s, 1H); *m/z* 451 (M+H)⁺, 449 (M-H)⁻

3-(3,5-Difluorophenoxy)-N-[4-chloromethyl-1,3-thiazol-2-yl]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]benzamide




To a stirred solution of 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{3,5-difluorophenoxy}benzoic acid (3.06mmol) in dichloromethane (20mL) was added 3 drops of DMF and oxalyl chloride (6.12mmol; 2.0 equivalents) dropwise, and the resulting mixture stirred at ambient temperature for 5 hours. The reaction mixture was concentrated *in vacuo*, azeotroped with toluene and dried overnight at reduced pressure. The residue was dissolved in dichloromethane and 4-chloromethylthiazol-2-ylamine (3.36mmol), triethylamine (3.36mmol) and dimethylaminopyridine (0.31mmol) added. The resulting mixture was stirred for 16 hours at ambient temperature. The reaction mixture was washed sequentially with 2M hydrochloric acid and 1M sodium hydrogencarbonate solution, dried (MgSO₄), and

concentrated *in vacuo*. The residue was chromatographed (eluting with 15-20% ethyl acetate in isohexane) to give the desired compound (33% yield). ^1H NMR δ (d_6 -DMSO): 1.24 (d, 3H), 3.28 (s, 3H obscured by solvent peak), 3.49 (m, 2H), 4.76 (m, 3H), 6.84 (dd, 2H), 6.94 (s, 1H), 7.04 (m, 1H), 7.32 (m, 2H), 7.55 (s, 1H), 12.77 (bs, 1H); m/z 469, 471 ($\text{M}+\text{H}$)⁺, 467, 5 469 ($\text{M}-\text{H}$)⁻


The synthesis of 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[3,5-difluorophenoxy]benzoic acid is described in Example 11 above.

10 **Example 14: 3-[(1S)-2-Methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]-N-1H-pyrazol-3-ylbenzamide**




Trifluoroacetic acid (0.5 mL) was added to a solution of *tert*-butyl 3-({3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]benzoyl} amino)-1H-pyrazole-1-carboxylate (180 mg, 0.330 mmol) in dry dichloromethane (3 mL) and the reaction was stirred under argon for 3 h. A further portion of trifluoroacetic acid (0.2 mL) was then added and the reaction was stirred for 30 min, then evaporated *in vacuo*. The residue was taken up in ethyl acetate (30 mL) and saturated aqueous sodium hydrogencarbonate (15 mL) and the residue was evaporated, then re-evaporated with dichloromethane/hexanes to produce the title 15 compound (145 mg, 100%) as a colourless foam; ^1H NMR δ ($\text{DMSO}-\text{d}_6$): 1.27 (d, 3H), 3.22 (s, 3H), 3.31 (s, 3H), 3.60 (m, 2H, partially obscured by HOD), 4.78 (m, 1H), 6.62 (s, 1H), 6.93 (s, 1H), 7.27 (d, 2H), 7.32 (s, 1H), 7.53 (s, 1H), 7.65 (s, 1H), 7.96 (d, 2H), 10.86 (s, 1H); m/z 444 ($\text{M}-\text{H}$)⁻

tert-Butyl 3-({3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]benzoyl}amino)-1H-pyrazole-1-carboxylate



O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (375 mg, 1.17 mmol) was added to 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenyl]oxy}benzoic acid (300 mg, 0.79 mmol) followed by addition of DMF (5 mL), diisopropylethylamine (0.35 mL) and *tert*-butyl 3-amino-1H-pyrazole-1-carboxylate (155 mg, 0.85 mmol). The reaction was stirred under argon for 4 h, the solvent evaporated, and the residue dissolved in saturated aqueous sodium hydrogencarbonate (30 mL) and ethyl acetate (50 mL). The organic layer was separated, washed with saturated aqueous ammonium chloride (30 mL), then dried (MgSO₄), filtered and evaporated. Purification by column chromatography eluting with 1:1 ethyl acetate:hexanes afforded the title compound (185 mg, 43%) as a colourless oil; ¹H NMR δ (CDCl₃): 1.37 (d, 3H), 1.63 (s, 9H), 3.09 (s, 3H), 3.40 (s, 3H), 3.58 (m, 2H), 4.61 (m, 1H), 6.85 (s, 1H), 7.08 (m, 2H), 7.15 (d, 2H), 7.30 (s, 1H), 7.92 (d, 2H), 8.01 (d, 1H), 8.58 (br. s, 1H); *m/z* 544 (M-H)⁺


tert-Butyl 3-amino-1H-pyrazole-1-carboxylate



1H-pyrazol-3-amine (428 mg, 5.15 mmol) was dissolved in dimethylformamide (5 mL) at 0°C and treated with sodium hydride (206 mg, 5.15 mmol) followed by stirring for a further 30 min. Warmed di-*tert*-butyl dicarbonate (1.12 g, 5.15 mmol) was then slowly added via syringe over 5 min and the reaction was allowed to warm to room temperature and stirred for a further 2 h. The reaction was taken up in saturated aqueous sodium hydrogencarbonate (50 mL) and ethyl acetate (100 mL). The organic layer was separated then dried (MgSO₄), filtered and evaporated. Purification by column chromatography (eluting with 1:1 ethyl acetate:hexanes to neat ethyl acetate) afforded the title compound (117 mg, 18%) as a white solid; ¹H NMR δ (CDCl₃): 1.62 (s, 9H), 4.00 (br. s, 2H), 5.81 (d, 1H), 7.82 (d, 1H)


The synthesis of 3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[[4-(methylsulfonyl)phenyl]oxy]benzoic acid is described in **Example 11** above.

Example 15: 3-[(1*S*)-2-Methoxy-(1-methylethyl)oxy]-*N*-(5-methyl-1*H*-pyrazol-3-yl)-5-[4-(methylsulfonyl)phenoxy]benzamide



Trifluoroacetic acid (1.5 mL) was added to a solution of *tert*-butyl 3-({3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]benzoyl}amino)-5-methyl-1*H*-pyrazole-1-carboxylate (500 mg, 0.330 mmol) in dry dichloromethane (6 mL) and the reaction was stirred under argon for 2 h. The solvent was removed in *vacuo* and the residue was taken up in ethyl acetate (30 mL) and saturated aqueous sodium hydrogencarbonate (15 mL). The organic layer was separated, dried (MgSO₄), filtered, evaporated, then re-evaporated with dichloromethane / hexanes to produce the title compound (350 mg, 85%) as a colourless foam; ¹H NMR δ (DMSO-*d*₆): 1.23 (d, 3H), 2.20 (s, 3H), 3.20 (s, 3H), 3.30 (s, 3H) (obscured by HOD), 3.50 (m, 2H) 4.78 (m, 1H), 6.38 (s, 1H), 6.90 (s, 1H), 7.22 (d, 2H), 7.30 (s, 1H), 7.45 (s, 1H), 7.93 (d, 2H), 10.71 (br. s, 1H), 12.08 (br. s, 1H); *m/z* 458 (M-H)⁺

***tert*-Butyl 3-({3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]benzoyl}amino)-5-methyl-1*H*-pyrazole-1-carboxylate**




O-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (500 mg, 1.31 mmol) was added to 3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[[4-(methylsulfonyl)phenyl]oxy]benzoic acid (400 mg, 1.05 mmol) followed by addition of DMF (6 mL),

diisopropylethylamine (0.47 mL) and *tert*-butyl 3-amino-5-methyl-1*H*-pyrazole-1-carboxylate (380 mg, 1.93 mmol). The reaction was stirred under argon for 72 h, then dissolved in saturated aqueous sodium hydrogencarbonate (30 mL) and ethyl acetate (50 mL). The organic layer was separated, washed with saturated aqueous ammonium chloride (30 mL), then dried (MgSO₄), filtered and evaporated. Purification by column chromatography eluting with 1:1 to 2:1 ethyl acetate:hexanes afforded the title compound (500 mg, 85%) as a foam; ¹H NMR δ (CDCl₃): 1.37 (d, 3H), 1.62 (s, 9H), 2.54 (s, 3H), 3.08 (s, 3H), 3.40 (s, 3H), 3.58 (m, 2H), 4.60 (m, 1H), 6.82 (m, 2H), 7.08 (m, 1H), 7.15 (d, 2H), 7.30 (s, 1H), 7.93 (d, 2H), 8.52 (br. s, 1H); *m/z* 558 (M-H)⁺

10

tert-Butyl 3-amino-5-methyl-1*H*-pyrazole-1-carboxylate




5-Methyl-1*H*-pyrazol-3-amine (800 mg, 8.25 mmol) was dissolved in dimethylformamide (10 mL) at 0 °C and treated with sodium hydride (336 mg, 8.25 mmol) followed by stirring for a further 30 min. Warmed di-*tert*-butyl dicarbonate (1.80 g, 8.25 mmol) was then slowly added via syringe over 5 min and the reaction was allowed to warm to room temperature and stirred for a further 1 h. The reaction was taken up in saturated aqueous sodium hydrogencarbonate (50 mL) and ethyl acetate (100 mL). The organic layer was separated then dried (MgSO₄), filtered and evaporated. Purification by column chromatography (eluting with 1:1 ethyl acetate:hexanes to 100% ethyl acetate) afforded the title compound (380 mg, 23%) as a colourless oil; ¹H NMR δ (CDCl₃): 1.62 (s, 9H), 2.43 (s, 3H), 3.87 (br. s, 2H), 5.60 (s, 1H)

20

The synthesis of 3-((1*S*)-2-methoxy-(1-methylethyl)oxy)-5-([4-(methylsulfonyl)phenyl]oxy)benzoic acid is described in **Example 11** above.


25

Example 16: 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-[4-(methoxymethyl)-1,3-thiazol-2-yl]-5-[4-(methylsulfonyl)phenoxy]benzamide



To a stirred solution of *N*-[4-(chloromethyl)-1,3-thiazol-2-yl]-3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]benzamide (280mg; 0.55mmol) in methanol (5mL) was added sodium methoxide (1.1mmol; 2.0 equiv; 25% weight in methanol) and the reaction mixture heated to 50°C and stirred overnight. The reaction mixture was concentrated *in vacuo* and chromatographed (eluting with 50-70% ethyl acetate:isohexane) to give the title compound (71mg; 26%); ¹H NMR δ (d₆-DMSO): 1.21 (d, 3H), 2.50 (2s, 6H, partially obscured by water peak), 3.21 (s, 3H), 3.43-3.54 (m, 2H), 4.40 (s, 2H), 4.78 (m, 1H), 6.98 (s, 1H), 7.12 (s, 1H), 7.25 (d, 2H), 7.37 (s, 1H), 7.58 (s, 1H), 7.95 (d, 2H), 12.69 (br s, 1H); *m/z* 507 (M+H)⁺, 505 (M-H)⁻

***N*-[4-(chloromethyl)-1,3-thiazol-2-yl]-3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenoxy]benzamide**




To a stirred solution of 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[4-(methylsulfonyl)phenyl]oxy}benzoic acid (1.0mmol) in dichloromethane (10mL) was added 1 drop of dimethylformamide and oxalyl chloride (2.0mmol; 2.0 equivalents) dropwise. The reaction mixture was stirred at ambient temperature under argon for two hours, then concentrated *in vacuo* and azeotroped with dichloromethane. The residue was dissolved in dichloromethane and 4-(chloromethyl)-1,3-thiazol-2-amine (*J. Indian Chem. Soc.* **1960**, *37*, 241; 1.0mmol) in dichloromethane was added along with diisopropylethylamine (2.5mmol) and dimethylaminopyridine (0.1 mmol). The resulting mixture was stirred for 13 hours under argon at ambient temperature, then concentrated *in vacuo* and chromatographed (eluting with 50-60% ethyl acetate in isohexane) to give the title compound (53% yield); ¹H NMR δ (d₆-

DMSO): 1.3 (d, 3H), 3.2 (s, 3H), 3.25 (s, 3H) 3.45 (m, 2H), 4.75 (s, 2H), 4.8 (m, 1H), 7.0 (s, 1H), 7.25 (d, 2H), 7.3 (s, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 7.95 (d, 2H), 12.80 (br s, 1H)


The synthesis of 3-((1S)-2-methoxy-(1-methylethyl)oxy)-5-({4-(methanesulfonyl)phenyl}oxy)benzoic acid is described in **Example 11** above.

Example 17: 3-[4-(azetidin-1-ylcarbonyl)phenoxy]-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide



- 10 To a suspension of 4-({3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[(3-methyl-1,2,4-thiadiazol-2-ylamino)carbonyl] phenyl}oxy)benzoic acid (300 mg), *O*-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (336 mg) and azetidine hydrochloride (190 mg) in DMF (5mL), was added diisopropylethylamine (0.68mL) and the mixture stirred at ambient temperature for 16 h. Water (75mL) was added and the mixture extracted with
- 15 ethyl acetate (3 x 25mL). The combined organic extracts were washed 1M aqueous hydrochloric acid (25 mL), saturated aqueous sodium hydrogen carbonate solution (25mL), brine, dried (MgSO₄), and evaporated to a residue which was chromatographed on silica with ethyl acetate as eluant to give the desired compound (190 mg). ¹H NMR δ (d₆-DMSO): 1.25 (d, 3H), 2.2-2.3 (m, 2H), 2.5 (m, 3H), 3.3 (s, 3H), 3.5 (m, 2H), 4.0 (m, 2H), 4.3 (m, 2H), 4.8 (m, 1H), 6.95 (s, 1H), 7.1 (d, 2H), 7.35 (s, 1H), 7.55 (s, 1H), 7.65 (d, 2H), 13.35 (s, 1H); *m/z*
- 20 483 (M+H)⁺


4-({3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[(3-methyl-1,2,4-thiadiazol-2-ylamino)carbonyl]phenyl}oxy)benzoic acid



A solution of ethyl 4-({3-{[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[(3-methyl-1,2,4-thiadiazol-2-ylamino)carbonyl] phenyl}oxy)benzoate (1.3g) in THF (40 mL) was added to a solution of lithium hydroxide monohydrate (310mg) in water (20mL). The mixture was stirred at ambient temperature for 16 hours and the THF removed *in vacuo*. The aqueous

- 5 layer was acidified with 1M hydrochloric acid (6.9mL), and the solid precipitate filtered off, washed with water and dried *in vacuo* to give the desired compound (1.12g). ^1H NMR δ (d_6 -DMSO): 1.2 (d, 3H), 2.45 (s, 3H), 3.25 (s, 3H), 3.5 (m, 2H), 4.7-4.8 (m, 1H), 6.95 (s, 1H), 7.1 (d, 2H), 7.35 (s, 1H), 7.6 (s, 1H), 7.95 (d, 2H); m/z 444 ($M+H$) $^+$

- 10 Ethyl 4-({3-{[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[(3-methyl-1,2,4-thiadiazol-2-ylamino)carbonyl] phenyl}oxy)benzoate




A solution of 3-hydroxy-5-{[(1*S*)-2-methoxy-(1-methylethyl)oxy]-*N*-(3-methyl-1,2,4-thiadiazol-2-yl)benzamide (3.23g), 4-ethoxycarbonylphenylboronic acid (3.63g), copper (II)

- 15 acetate (3.63g), triethylamine (6.9mL) and freshly activated 4Å molecular sieves (12.5g) in dichloromethane (250mL) was stirred at ambient temperature and under ambient atmosphere for 2 days. The reaction mixture was filtered through diatomaceous earth, washed with dichloromethane (2 x 50mL), the dichloromethane removed *in vacuo* and the residual oil partitioned between ethyl acetate (300mL) and 1M hydrochloric acid (200mL). The ethyl

- 20 acetate layer was separated, washed sequentially with aqueous sodium hydrogen carbonate solution and brine, dried (MgSO_4), and evaporated to a residue which was chromatographed on silica with 40% ethyl acetate in isohexane as eluant to give the desired compound (1.35g).


- ^1H NMR δ (CDCl_3): 1.3 (d, 3H), 1.4 (t, 3H), 2.45 (s, 3H), 3.4 (s, 3H), 3.5-3.6 (m, 2H), 4.35 (q, 2H), 4.5-4.6 (m, 1H), 6.85 (s, 1H), 7.0 (d, 2H), 7.1 (s, 1H), 7.3 (d, 1H), 8.05 (d, 2H), 10.5 (s, 1H); m/z 472 ($M+H$) $^+$
- 25

3-Hydroxy-5-{[(1S)-2-methoxy-(1-methylethyl)oxy]}-N-(3-methyl-1,2,4-thiadiazol-2-yl)benzamide



A solution of 3-{[(1S)-2-methoxy-(1-methylethyl)oxy]}-5-{phenylmethoxy}-N-(3-methyl-1,2,4-thiadiazol-2-yl)benzamide (9.53g) and thioanisole (13.9mL) in trifluoroacetic acid (45mL) was stirred at ambient temperature for 16 hours. The trifluoroacetic acid was removed *in vacuo* and the residual oil partitioned between ethyl acetate (100mL) and aqueous sodium hydrogen carbonate solution (300mL). The aqueous layer was separated, extracted with ethyl acetate (2 x 100mL), and the combined organic extracts washed with brine, dried (MgSO₄), and evaporated to a residue which was chromatographed on silica with 50% ethyl acetate in isohexane as eluant to give the desired compound (4.5g). ¹H NMR δ (CDCl₃): 1.2 (d, 3H), 2.5 (s, 3H), 3.3 (s, 3H), 3.4-3.6 (m, 2H), 4.6-4.7 (m, 1H), 6.6 (s, 1H), 7.05 (s, 1H), 7.1 (s, 1H), 9.85 (s, 1H), 13.2 (s, 1H); *m/z* 324 (M+H)⁺

3-{[(1S)-2-Methoxy-(1-methylethyl)oxy]}-5-{phenylmethoxy}-N-(3-methyl-1,2,4-thiadiazol-2-yl)benzamide




To a solution of 3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-{[phenylmethyl]oxy}benzoic acid (15.8g) in dichloromethane (260mL) was added oxalyl chloride (5.24mL), followed by DMF (1 drop), and the mixture stirred at ambient temperature for 16 hours. The dichloromethane and excess oxalyl chloride were removed *in vacuo*, the residual oil dissolved in dichloromethane (50mL) and added to a solution of 5-amino-3-methyl-1,2,4 thiadiazole (6.05g) and triethylamine (14.6mL) in dichloromethane (150mL) at 0-5°C, and the mixture stirred at ambient temperature for 16 hours. The dichloromethane and excess triethylamine were removed *in vacuo*, and the residual oil partitioned between ethyl acetate (250mL) and 1M hydrochloric acid (150mL). The ethyl acetate layer was separated, washed sequentially

with 1M hydrochloric acid, aqueous sodium hydrogen carbonate solution, and brine, dried (MgSO₄), and evaporated to a residue which was chromatographed on alumina with ethyl acetate as eluant, then on silica with 30% ethyl acetate in isohexane as eluant to give the desired compound (9.6g). ¹H NMR δ (CDCl₃): 1.3 (d, 3H), 2.45 (s, 3H), 3.4 (s, 3H), 3.5-3.6 (m, 2H), 4.55-4.6 (m, 1H), 5.05 (s, 2H), 6.8 (s, 1H), 7.1 (m, 2H), 7.25 (m, 5H), 10.7 (s, 1H); *m/z* 414 (M+H)⁺

The synthesis of 3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-{[phenylmethyl]oxy}benzoic acid is described in **Example 3** above.

10


Example 18: 3-[(1*S*)-2-Methoxy-(1-methylethyl)oxy]-5-(4-[(1-methylpiperidin-4-yl)amino]carbonyl]phenoxy)-*N*-(3-methyl-1,2,4-thiadiazol-5-yl)benzamide



In a similar manner to that described above for **Example 17**, **Example 18** was prepared using 4-({3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[(3-methyl-1,2,4-thiadiazol-2-ylamino)carbonyl]phenyl}oxy)benzoic acid and the appropriate amine. ¹H NMR δ (d₆-DMSO): 1.25 (d, 3H), 1.5-1.65 (m, 2H), 1.7-1.8 (m, 2H), 1.95-2.0 (m, 2H), 2.15 (s, 3H), 2.3 (s, 3H), 2.7-2.8 (m, 2H), 3.3 (s, 3H), 3.55 (m, 2H), 3.7 (m, 1H), 4.7 (m, 1H), 6.7 (s, 1H), 7.1 (d, 2H), 7.25 (s, 1H), 7.5 (s, 1H), 7.85 (d, 2H), 8.2 (d, 1H); *m/z* 540 (M+H)⁺

20

Example 19: 3-[4-(Azetidin-1-ylcarbonyl)-2-chlorophenoxy]-5-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-*N*-(1-methyl-1*H*-pyrazol-3-yl)benzamide






To a suspension of 3-chloro-4-[(3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[(1-methyl-1*H*-pyrazol-3-yl)amino]carbonyl]phenoxy]benzoic acid (344 mg), *O*-(7-azabenzotriazol-1-yl)-

25

1,1,3,3-tetramethyluronium hexafluorophosphate (366 mg) and azetidine hydrochloride (88 mg) in DMF (10mL), was added diisopropylethylamine (0.50mL) and the mixture stirred at ambient temperature for 24 hours. Water (30mL) was added and the mixture extracted with ethyl acetate (3 x 15mL). The combined organic extracts were washed with brine, dried (MgSO₄), and evaporated to a residue which was chromatographed on silica eluting with a gradient of 50-100% ethyl acetate in hexane to give the desired compound (197 mg). ¹H NMR δ (CDCl₃): 1.3 (d, 3H), 2.4 (m, 2H), 3.4 (s, 3H), 3.5 (m, 2H), 3.8 (s, 3H), 4.2-4.4 (m, 4H), 4.6 (m, 1H), 6.7 (d, 2H), 7.0 (m, 2H), 7.2 (m, 2H), 7.5 (d, 1H), 7.8 (d, 1H), 8.60 (br s, 1H); *m/z* 499 (M+H)⁺


10

In a similar manner, Examples 19a-19c were also prepared:-

Example	Structure	<i>m/z</i>	NMR
19a		483 (M+H) ⁺	¹ H NMR δ (CDCl ₃): 1.3 (d, 3H), 2.4 (m, 2H), 3.4 (s, 3H), 3.5 (m, 2H), 3.8 (s, 3H), 4.2-4.4 (m, 4H), 4.6 (m, 1H), 6.8 (m, 2H), 7.0 (m, 2H), 7.2 (m, 1H), 7.3 (m, 1H), 7.4 (d, 1H), 7.5 (d, 1H), 8.50 (s, 1H)
19b		533 (M+H) ⁺	¹ H NMR δ (CDCl ₃): 1.3 (d, 3H), 2.4 (m, 2H), 3.4 (s, 3H), 3.5 (m, 2H), 3.8 (s, 3H), 4.2-4.4 (m, 4H), 4.6 (m, 1H), 6.8 (m, 2H), 6.95 (d, 1H), 7.1 (s, 1H), 7.3 (m, 2H), 7.75 (d, 1H), 8.0 (s, 1H), 8.50 (s, 1H)
19c		453 (M+H) ⁺ 451 (M-H) ⁻	¹ H NMR δ (DMSO-d ₆): 1.2 (d, 3H), 2.95 (s, 6H), 3.25 (s, 3H), 3.5 (m, 2H), 3.8 (s, 3H), 4.75 (m, 1H), 6.55 (s, 1H), 6.8 (s, 1H), 7.05 (d, 2H), 7.2 (s, 1H), 7.4 (s, 1H), 7.45 (d, 2H), 7.6 (s, 1H), 10.85 (br s, 1H)

The required acids for Examples 19 & 19a-19c were prepared as described below:-

3-Chloro-4-[(3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[(1-methyl-1H-pyrazol-3-yl)amino]carbonyl}phenyl)oxy]benzoic acid




To a solution of methyl 3-chloro-4-(3-[(1S)-2-methoxy-(1-methylethyl)oxy]-5-[(1-methyl-1H-pyrazol-3-yl)amino]carbonyl}phenoxy)benzoate (2.23g) in THF (58 mL) was added a 1M solution of lithium hydroxide monohydrate in water (11.7mL). The mixture was stirred at ambient temperature for 18 hours and the THF removed *in vacuo*. The aqueous layer was acidified with 2M hydrochloric acid (5.85mL), and the solid precipitate filtered off, washed with water and dried *in vacuo* to give the desired acid (1.87g). ¹H NMR δ (CDCl₃): 1.4 (d, 3H), 3.4 (s, 3H), 3.6 (m, 2H), 3.8 (s, 3H), 4.7 (m, 1H), 6.95 (m, 1H), 7.05 (m, 1H), 7.1 (d, 1H), 7.3 (m, 2H), 7.6 (m, 2H), 8.1 (s, 1H), 10.75 (br s, 1H); *m/z* 460 (M+H)⁺

The acids required for the synthesis of **Examples 19a-19c** were made using an analogous method:

Structure	<i>m/z</i>	NMR
	444 (M+H) ⁺	¹ H NMR δ (CDCl ₃): 1.4 (d, 3H), 3.4 (s, 3H), 3.5 (m, 2H), 3.8 (s, 3H), 4.7 (m, 1H), 6.95 (m, 1H), 7.0 (m, 1H), 7.1 (m, 1H), 7.3 (m, 2H), 7.6 (m, 2H), 7.8 (d, 1H), 10.8 (s, 1H)
	494 (M+H) ⁺	¹ H NMR δ (d ₆ DMSO): 1.2 (d, 3H), 3.3 (s, 3H), 3.5 (m, 2H), 3.8 (s, 3H), 4.7 (m, 1H), 6.5 (s, 1H), 6.8 (s, 1H), 7.05 (d, 1H), 7.2 (s, 1H), 7.5 (s, 1H), 7.6 (s, 1H), 8.1(d, 1H), 8.5 (s, 1H), 10.85 (s, 1H)
	426 (M+H) ⁺ 424 (M-H) ⁻	¹ H NMR δ (d ₆ DMSO): 1.2 (d, 3H), 3.25 (s, 3H), 3.5 (m, 2H), 3.8 (s, 3H), 4.75 (m, 1H), 6.55 (s, 1H), 6.85 (s, 1H), 7.1 (d, 2H), 7.2 (s, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.0(d, 2H), 10.85 (br s, 1H)

The required esters for **Examples 19 & 19a-19c** were prepared as described below:-


Methyl 3-chloro-4-(3-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-5-[(1-methyl-1*H*-pyrazol-3-yl)amino]carbonyl}phenoxy)benzoate



- 5 To a solution of 3-hydroxy-5-[(1*S*)-2-methoxy-(1-methylethyl)oxy]-*N*-(1-methyl-1*H*-pyrazol-3-yl)benzamide (832mg, 2.72mmol) and methyl-3-chloro-4-fluorobenzoate (504 mg, 2.72mmol) in acetonitrile (20 mL) was added potassium carbonate (364 mg, 2.72mmol) and the stirred mixture heated at 160°C in a 'Smith Creator Microwave' for 30 minutes. The mixture was allowed to return to ambient temperature and pressure, filtered and evaporated to
- 10 a residue which was chromatographed on silica with 0 -50% ethyl acetate in hexane as eluant to give the desired compound (1.11g); m/z 474 ($M+H$)⁺

The esters required for the synthesis of **Examples 19a-19c** were prepared using an analogous method:

Structure ^s	m/z	NMR
	458 ($M+H$) ⁺	¹ H NMR δ (CDCl ₃): 1.3(d, 3H), 3.4 (s, 3H), 3.5 (m, 2H), 3.8 (s, 3H), 3.9 (s, 3H), 4.6 (m, 1H), 6.9 (m, 2H), 7.1 (m, 2H), 7.3 (m, 2H), 7.8 (m, 2H), 8.6 (s, 1H)
	508 ($M+H$) ⁺	¹ H NMR δ (CDCl ₃): 1.3 (d, 3H), 3.4 (s, 3H), 3.5 (m, 2H), 3.8 (s, 3H), 3.95 (s, 3H), 4.6 (m, 1H), 6.75 (d, 1H), 6.85 (m, 1H), 6.95 (d, 1H), 7.1 (m, 1H), 7.3 (m, 1H), 7.35(m, 1H), 8.1(d, 1H), 8.4 (s, 1H), 8.5 (s, 1H)

	454 (M+H) ⁺	¹ H NMR δ (d ₆ DMSO): 1.2 (d, 3H), 1.3 (t, 3H), 3.25 (s, 3H), 3.5 (m, 2H), 3.8 (s, 3H), 4.3 (q, 2H), 4.8 (m, 1H), 6.55 (s, 1H), 6.8 (s, 1H), 7.1 (d, 2H), 7.2 (s, 1H), 7.45 (s, 1H), 7.6 (s, 1H), 8.0 (d, 2H), 10.85 (br s, 1H)
---	------------------------	--

^sThe precursor for **Example 19c** was prepared at 150°C in DMF for 4 hours, using 1.2 equivalents of the fluoro-ester

The synthesis of 3-hydroxy-5-[(1S)-2-methoxy-(1-methylethyl)oxy]-N-(1-methyl-1H-pyrazol-3-yl)benzamide is described in **Example 3**.

BIOLOGICAL

Tests:

The biological effects of the compounds of formula (I) may be tested in the following way:

(1) Enzymatic activity

Enzymatic activity of recombinant human pancreatic GLK may be measured by incubating GLK, ATP and glucose. The rate of product formation may be determined by coupling the assay to a G-6-P dehydrogenase, NADP/NADPH system and measuring the increase in optical density at 340nm (Matschinsky et al 1993).

Production of recombinant GLK and GLKRP:

Human GLK and GLKRP cDNA was obtained by PCR from human pancreatic and hepatic mRNA respectively, using established techniques described in Sambrook J, Fritsch EF & Maniatis T, 1989. PCR primers were designed according to the GLK and GLKRP cDNA sequences shown in Tanizawa et al 1991 and Bonthron, D.T. *et al* 1994 (later corrected in Warner, J.P. 1995).

Cloning in Bluescript II vectors

GLK and GLKRP cDNA was cloned in *E. coli* using pBluescript II, (Short et al 1998) a recombinant cloning vector system similar to that employed by Yanisch-Perron C *et al* (1985), comprising a *colEI*-based replicon bearing a polylinker DNA fragment containing

multiple unique restriction sites, flanked by bacteriophage T3 and T7 promoter sequences; a filamentous phage origin of replication and an ampicillin drug resistance marker gene.

Transformations

5 E. Coli transformations were generally carried out by electroporation. 400 mL cultures of strains DH5a or BL21(DE3) were grown in L-broth to an OD 600 of 0.5 and harvested by centrifugation at 2,000g. The cells were washed twice in ice-cold deionised water, resuspended in 1mL 10% glycerol and stored in aliquots at -70°C . Ligation mixes were desalted using Millipore V series™ membranes (0.0025mm) pore size). 40mL of cells were
10 incubated with 1mL of ligation mix or plasmid DNA on ice for 10 minutes in 0.2cm electroporation cuvettes, and then pulsed using a Gene Pulser™ apparatus (BioRad) at 0.5kVcm^{-1} , 250mF. Transformants were selected on L-agar supplemented with tetracycline at 10mg/mL or ampicillin at 100mg/mL.

15 *Expression*

GLK was expressed from the vector pTB375NBSE in E.coli BL21 cells,, producing a recombinant protein containing a 6-His tag immediately adjacent to the N-terminal methionine. Alternatively, another suitable vector is pET21(+)DNA, Novagen, Cat number 697703. The 6-His tag was used to allow purification of the recombinant protein on a column

20 packed with nickel-nitrilotriacetic acid agarose purchased from Qiagen (cat no 30250).

GLKRP was expressed from the vector pFLAG CTC (IBI Kodak) in E.coli BL21 cells, producing a recombinant protein containing a C-terminal FLAG tag. The protein was purified initially by DEAE Sepharose ion exchange followed by utilisation of the FLAG tag for final purification on an M2 anti-FLAG immunoaffinity column purchased from Sigma-Aldrich (cat

25 no. A1205).



(2) Oral Glucose Tolerance Test (OGTT)

Oral glucose tolerance tests were done on conscious Zucker obese fa/fa rats (age 12-13 weeks or older) fed a high fat diet (45 % kcal fat) for at least two weeks prior to

30 experimentation. The animals were fasted for 2 hours before use for experiments. A test compound or a vehicle was given orally 120 minutes before oral administration of a glucose solution at a dose of 2 g/kg body weight. Blood glucose levels were measured using a

Accucheck glucometer from tail bled samples taken at different time points before and after administration of glucose (time course of 60 minutes). A time curve of the blood glucose levels was generated and the area-under-the-curve (AUC) for 120 minutes was calculated (the time of glucose administration being time zero). Percent inhibition was determined using the

5 AUC in the vehicle-control group as zero percent inhibition.

**Example 11b****Example II107**

Compounds of the invention generally have an activating activity for glucokinase with an EC_{50} of less than about 500nM. For example, Example 11b has an EC_{50} of 30nM.

10 Example 11b and Example II107 in WO 03/015774 have broadly similar EC_{50} values. However Example 11b has superior oral exposure and exhibits 29% OGTT activity at 10 mg/kg but Example II107 in WO 03/015774 is not active at 10 mg/kg.

REFERENCES


- 15 1 Printz, R. L., Magnuson, M. A. and Granner, D. K. (1993) Annual Review of Nutrition **13**, 463-96
- 2 DeFronzo, R. A. (1988) Diabetes **37**, 667-87
- 3 Froguel, P., Zouali, H., Vionnet, N., Velho, G., Vaxillaire, M., Sun, F., Lesage, S., Stoffel, M., Takeda, J. and Passa, P. (1993) New England Journal of Medicine **328**, 697-702
- 20 4 Bell, G. I., Pilkis, S. J., Weber, I. T. and Polonsky, K. S. (1996) Annual Review of Physiology **58**, 171-86
- 5 Velho, G., Petersen, K. F., Perseghin, G., Hwang, J. H., Rothman, D. L., Pueyo, M. E., Cline, G. W., Froguel, P. and Shulman, G. I. (1996) Journal of Clinical Investigation **98**, 1755-61
- 25 6 Christesen, H. B., Jacobsen, B. B., Odili, S., Buettger, C., Cuesta-Munoz, A., Hansen, T., Brusgaard, K., Massa, O., Magnuson, M. A., Shiota, C., Matschinsky, F. M. and Barbetti, F. (2002) Diabetes **51**, 1240-6

- 6a Gloyn, A.L., Noordam, K., Willemsen, M.A.A.P., Ellard, S., Lam, W.W.K., Campbell, I. W., Midgley, P., Shiota, C., Buettger, C., Magnuson, M.A., Matschinsky, F.M., and Hattersley, A.T.; *Diabetes* **52**: 2433-2440
- 7 Glaser, B., Kesavan, P., Heyman, M., Davis, E., Cuesta, A., Buchs, A., Stanley, C. A., Thornton, P. S., Permutt, M. A., Matschinsky, F. M. and Herold, K. C. (1998) *New England Journal of Medicine* **338**, 226-30
- 8 Caro, J. F., Triester, S., Patel, V. K., Tapscott, E. B., Frazier, N. L. and Dohm, G. L. (1995) *Hormone & Metabolic Research* **27**, 19-22
- 9 Desai, U. J., Slosberg, E. D., Boettcher, B. R., Caplan, S. L., Fanelli, B., Stephan, Z., Gunther, V. J., Kaleko, M. and Connelly, S. (2001) *Diabetes* **50**, 2287-95
- 10 Shiota, M., Postic, C., Fujimoto, Y., Jetton, T. L., Dixon, K., Pan, D., Grimsby, J., Grippo, J. F., Magnuson, M. A. and Cherrington, A. D. (2001) *Diabetes* **50**, 622-9
- 11 Ferre, T., Pujol, A., Riu, E., Bosch, F. and Valera, A. (1996) *Proceedings of the National Academy of Sciences of the United States of America* **93**, 7225-30
- 15 12 Seoane, J., Barbera, A., Telemaque-Potts, S., Newgard, C. B. and Guinovart, J. J. (1999) *Journal of Biological Chemistry* **274**, 31833-8
- 13 Moore, M. C., Davis, S. N., Mann, S. L. and Cherrington, A. D. (2001) *Diabetes Care* **24**, 1882-7
- 14 Alvarez, E., Roncero, I., Chowen, J. A., Vazquez, P. and Blazquez, E. (2002) *Journal of Neurochemistry* **80**, 45-53
- 20 15 Lynch, R. M., Tompkins, L. S., Brooks, H. L., Dunn-Meynell, A. A. and Levin, B. E. (2000) *Diabetes* **49**, 693-700
- 16 Roncero, I., Alvarez, E., Vazquez, P. and Blazquez, E. (2000) *Journal of Neurochemistry* **74**, 1848-57
- 25 17 Yang, X. J., Kow, L. M., Funabashi, T. and Mobbs, C. V. (1999) *Diabetes* **48**, 1763-1772
- 18 Schuit, F. C., Huypens, P., Heimberg, H. and Pipeleers, D. G. (2001) *Diabetes* **50**, 1-11
- 19 Levin, B. E. (2001) *International Journal of Obesity* **25**, supp 5, S68-S72
- 20 Alvarez, E., Roncero, I., Chowen, J. A., Thorens, B. and Blazquez, E. (1996) *Journal of Neurochemistry* **66**, 920-7
- 30 21 Mobbs, C. V., Kow, L. M. and Yang, X. J. (2001) *American Journal of Physiology - Endocrinology & Metabolism* **281**, E649-54

- 22 Levin, B. E., Dunn-Meynell, A. A. and Routh, V. H. (1999) *American Journal of Physiology* **276**, R1223-31
- 23 Spanswick, D., Smith, M. A., Groppi, V. E., Logan, S. D. and Ashford, M. L. (1997) *Nature* **390**, 521-5
- 5 24 Spanswick, D., Smith, M. A., Mirshamsi, S., Routh, V. H. and Ashford, M. L. (2000) *Nature Neuroscience* **3**, 757-8
- 25 Levin, B. E. and Dunn-Meynell, A. A. (1997) *Brain Research* **776**, 146-53
- 26 Levin, B. E., Govek, E. K. and Dunn-Meynell, A. A. (1998) *Brain Research* **808**, 317-9
- 27 Levin, B. E., Brown, K. L. and Dunn-Meynell, A. A. (1996) *Brain Research* **739**, 293-
10 300
- 28 Rowe, I. C., Boden, P. R. and Ashford, M. L. (1996) *Journal of Physiology* **497**, 365-77
- 29 Fujimoto, K., Sakata, T., Arase, K., Kurata, K., Okabe, Y. and Shiraishi, T. (1985) *Life Sciences* **37**, 2475-82
- 30 Kurata, K., Fujimoto, K. and Sakata, T. (1989) *Metabolism: Clinical & Experimental*
15 **38**, 46-51
- 31 Kurata, K., Fujimoto, K., Sakata, T., Etou, H. and Fukagawa, K. (1986) *Physiology & Behavior* **37**, 615-20
- 32 Jetton T.L., Liang Y., Pettepher C.C., Zimmerman E.C., Cox F.G., Horvath K., Matschinsky F.M., and Magnuson M.A., *J. Biol. Chem.*, Feb **1994**; **269**: 3641 - 3654 .
- 20 33 Reimann F. and Gribble F. M., *Diabetes* **2002** **51**: 2757-2763
- 34 Cheung A. T., Dayanandan B., Lewis J. T., Korbitt G. S., Rajotte R. V., Bryer-Ash M., Boylan M. O., Wolfe M. M., Kieffer T. J., *Science*, Vol 290, Issue 5498, 1959-1962 , 8 December 2000

Claims:

1. A compound of Formula (I):



wherein:

R^1 is methoxymethyl;

R^2 is selected from $-C(O)NR^4R^5$, $-SO_2NR^4R^5$, $-S(O)_pR^4$ and HET-2;

- 10 HET-1 is a 5- or 6-membered, C-linked heteroaryl ring containing a nitrogen atom in the 2-position and optionally 1 or 2 further ring heteroatoms independently selected from O, N and S; which ring is optionally substituted on an available carbon atom, or on a ring nitrogen atom provided it is not thereby quaternised, with 1 or 2 substituents independently selected from R^6 ;

- 15 HET-2 is a 4-, 5- or 6-membered, C- or N-linked heterocyclyl ring containing 1, 2, 3 or 4 heteroatoms independently selected from O, N and S, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$, and wherein a sulphur atom in the heterocyclic ring may optionally be oxidised to an $S(O)$ or $S(O)_2$ group, which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^7 ;

- 20 R^3 is selected from halo, fluoromethyl, difluoromethyl, trifluoromethyl, methyl, methoxy and cyano;

R^4 is selected from hydrogen, (1-4C)alkyl [optionally substituted by 1 or 2 substituents independently selected from HET-2, $-OR^5$, $-SO_2R^5$, (3-6C)cycloalkyl (optionally substituted with 1 group selected from R^7) and $-C(O)NR^5R^5$] and HET-2;

- 25 R^5 is hydrogen or (1-4C)alkyl;

or R^4 and R^5 together with the nitrogen atom to which they are attached may form a heterocyclyl ring system as defined by HET-3;

R^6 is independently selected from (1-4C)alkyl, halo, hydroxy(1-4C)alkyl, (1-4C)alkoxy(1-4C)alkyl, (1-4C)alkylS(O)p(1-4C)alkyl, amino(1-4C)alkyl, (1-4C)alkylamino(1-4C)alkyl, di(1-4C)alkylamino(1-4C)alkyl and HET-4;

R^7 is selected from $-OR^5$, (1-4C)alkyl, $-C(O)(1-4C)alkyl$, $-C(O)NR^4R^5$, (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl and $-S(O)pR^5$;

HET-3 is an N-linked, 4 to 6 membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 or 2 further heteroatoms (in addition to the linking N atom) independently selected from O, N and S, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$ and wherein a sulphur atom in the ring may optionally be oxidised to an S(O) or S(O)₂ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^8 ; or

HET-3 is an N-linked, 7 membered, saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further heteroatom (in addition to the linking N atom) independently selected from O, S and N, wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$ group and wherein a sulphur atom in the ring may optionally be oxidised to an S(O) or S(O)₂ group; which ring is optionally substituted on an available carbon or nitrogen atom by 1 or 2 substituents independently selected from R^8 ; or

HET-3 is an 6-10 membered bicyclic saturated or partially unsaturated heterocyclyl ring, optionally containing 1 further nitrogen atom (in addition to the linking N atom), wherein a $-CH_2-$ group can optionally be replaced by a $-C(O)-$; which ring is optionally substituted on an available carbon or nitrogen atom by 1 substituent selected from R^3 ;

R^8 is selected from $-OR^5$, (1-4C)alkyl, $-C(O)(1-4C)alkyl$, $-C(O)NR^4R^5$, (1-4C)alkylamino, di(1-4C)alkylamino, HET-3 (wherein said ring is unsubstituted), (1-4C)alkoxy(1-4C)alkyl, hydroxy(1-4C)alkyl and $-S(O)pR^5$;

HET-4 is a 5- or 6-membered, C- or N- linked unsubstituted heteroaryl ring containing 1, 2 or 3 ring heteroatoms independently selected from O, N and S;

p is (independently at each occurrence) 0, 1 or 2;



m is 0 or 1;

n is 0, 1 or 2;

provided that when m is 0, then n is 1 or 2;


or a salt, pro-drug or solvate thereof.

2. A compound of the formula (I) as claimed in Claim 1 wherein R¹ has the (S) configuration.
3. A compound of the formula (I) as claimed in Claim 1 or Claim 2, or a salt, pro-drug or solvate thereof, wherein HET-1 is a 5-membered ring.
4. A compound of the formula (I) as claimed in Claim 1, Claim 2, or Claim 3, or a salt, pro-drug or solvate thereof, wherein HET-3 is a 4- to 6-membered ring.
5. A pharmaceutical composition comprising a compound according to any one of Claims 1 to 4, or a salt, pro-drug or solvate thereof, together with a pharmaceutically acceptable diluent or carrier.
6. A compound according to any one of Claims 1 to 4 for use in the preparation of a medicament for treatment of a disease mediated through GLK.
7. A method of treating GLK mediated diseases by administering an effective amount of a compound of Formula (I) as claimed in any one of Claims 1 to 4 or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.
8. A process for the preparation of a compound of Formula (I) as claimed in any one of Claims 1 to 4, which comprises (wherein variables are as defined in Claim 1 unless otherwise stated):
- (a) reaction of an acid of Formula (III) or activated derivative thereof with a compound of Formula (IV),




or

- (b) reaction of a compound of Formula (V) with a compound of Formula (VI),




(V)




(VI)

wherein X^1 is a leaving group and X^2 is a hydroxyl group or X^1 is a hydroxyl group and X^2 is a leaving group;

- 5 [or by reaction with the intermediate ester Formula (VII), wherein P^1 is a protecting group followed by ester hydrolysis and amide formation];




(V)




(VII)

or

- 10 (c) reaction of a compound of Formula (VIII) with a compound of Formula (IX)




(VIII)




(IX)

wherein X^3 is a leaving group or an organometallic reagent and X^4 is a hydroxyl group or X^3 is a hydroxyl group and X^4 is a leaving group or an organometallic reagent;

- 15 [or by reaction or (VIII) with the intermediate ester Formula (X), followed by ester hydrolysis and amide formation];




(VIII)




(X)

or

(d) reaction of a compound of Formula (XI) with a compound of Formula (XII),



(XI)



(XII);

wherein X^5 is a leaving group;

5 and thereafter, if necessary:

i) converting a compound of Formula (I) into another compound of Formula (I);

ii) removing any protecting groups; and/or

iii) forming a salt, pro-drug or solvate.

PCT/GB2005/000545

